

OPTIMALITY INDEX – US

August 2011

REFERENCE CITATIONS

(The evidence in support of the criterion of optimality established for each item in the index)

Aagaard-Tillery KM, Porter TF, Lane RH, Varner MW, Lacoursiere DY. In utero tobacco exposure is associated with modified effects of maternal factors on fetal growth. *Am J Obstet Gynecol.* 2008;198(1):66.e1-6.

Abalos E, Duley L, Steyn DW, Henderson-Smart DJ. Antihypertensive drug therapy for mild to moderate hypertension during pregnancy. *Cochrane Database Syst Rev.* 2007 Jan 24(1):CD002252.

Abrao KC, Francisco RPV, Miyandahira S, Cicarelli DD, Zugaib M. Elevation of uterine basal tone and fetal health rate abnormalities after labor analgesia *Obstet Gynecol.* 2009; 113(1):41-47.

ACOG Committee on Practice Bulletins. Pregestational diabetes mellitus. *Clinical Management Guidelines for Obstetrician-Gynecologists.* Number 60, March 2005.

ACOG. Prevention of early-onset group B streptococcal disease in newborns. *ACOG Committee Opinion #279.* 2002.

Aebi-Popp K, Lapaire O, Glass TR, Vilen L, Rudin C, Elzi L, Battagay M, Keiser O, Martinez de Tejada B, Hoesli I. Pregnancy and delivery outcomes of HIV infected women in Switzerland 2003 – 2008. *J Perinat Med.* 2010. DOI:10.1515/JPM.2010.052

Alfirevic Z. Early amniocentesis versus transabdominal chorion villus sampling for prenatal diagnosis. In: *The Cochrane Database of Systematic Reviews,* 2003.

Alfirevic Z, Gosden CM, Neilson JP. Chorion villus sampling versus amniocentesis for prenatal diagnosis In: *The Cochrane Database of Systematic Reviews,* 2003.

Alfirevic Z, Neilson JP. Biophysical profile for fetal assessment in high risk pregnancies. In: *The Cochrane Database of Systematic Reviews,* 2003.

Algert C, Morris J, Simpson J, Ford J, Roberts C. Labor before a primary cesarean delivery. *Obstetrics & Gynecology.* 2008;112:1061-6.

Allen VM, Armson BA, Wilson RD, Allen VM, Blight C, Gagnon A, Johnson JA, Langlois S, Summers A, Wyatt P, Farine D, Armson BA, Crane J, Delisle MF, Keenan-Lindsay L, Morin V, Schneider CE, Van Aerde J. Teratogenicity associated with pre-existing and gestational diabetes. *J Obstet Gynaecol Can.* 2007; 29(11):927-44.

- Aliyu MH, Wilson RE, Zoorob R, Chakrabarty S, Alio AP, Kirby RS, Salihu HM. Alcohol consumption during pregnancy and the risk of early stillbirth among singletons. *Alcohol*. 2008;42(5):369-74.
- Althaus J, Petersen S, Fox H, Holcroft C, Graham E. Can electronic fetal monitoring identify preterm neonates with cerebral white matter injury? *Obstet Gynecol*. 2005;105(3):458-465.
- Alwan N, Tuffnell DJ, West J. Treatments for gestational diabetes. *Cochrane Database Syst Rev*. 2009 Jul 8(3):CD003395.
- Al-Zirqi I, Vangen S, Forsen L, Stray-Pedersen B. Prevalence and risk factors of severe obstetric haemorrhage. *BJOG*. 2008; 115(10):1265-72.
- American Academy of Pediatrics (AAP), ACOG. Use and abuse of the Apgar score. ACOG Committee Opinion 333. Washington, DC, 2006.
- Ananth CV, Wilcox AJ. Placental abruption and perinatal mortality in the United States. *Am J Epidemiol*. 2001;153(4):332-7.
- Anderson GC, Moore E, Hepworth J, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. In: *The Cochrane Database of Systematic Reviews*, 2003.
- Andreoli SP. Acute renal failure in the newborn. *Semin Perinatol*. 2004;28(2):112-23.
- Anim-Somuah M, Smyth RMD, Howell CJ. Epidural versus non-epidural or analgesia in labour (review). *Cochrane Database of Systematic Reviews*. 2005.
- Archabald KL, Friedman A, Raker CA, Anderson BL. Impact of trimester on morbidity of acute pyelonephritis in pregnancy. *Am J Obstet Gynecol*. 2009;201(4):406.e1-4. Epub2009 Aug 18.
- Arlettaz R, Kashiwagi M, Das-Kundu S, Fauchere JC, Lang A, Bucher JU. Methadone maintenance program in pregnancy in a Swiss perinatal center (II): neonatal outcome and social resources. *Acta Obstet Gynecol Scand*. 2005;84(2):145-50.
- Auger N, Daniel M, Platt RW, Luo ZC, Wu Y, Choinière R. The joint influence of marital status, interpregnancy interval, and neighborhood on small for gestational age birth: a retrospective cohort study. *BMC Childbirth*. 2008 Feb 28; 8:7
- Aziz N, Cheng YW, Caughey AB. Neonatal outcomes in the setting of preterm premature rupture of membranes complicated by chorioamnionitis. *J Matern Fetal Neonatal Med*. 2009; 22(9):780-4.
- Bada HS, Das A, Bauer CR, Shankaran S, Lester B, LaGasse L, Hammond J, Wright LL, Higgins R. Impact of prenatal cocaine exposure on child behavior problems through school age. *Pediatrics*. 2007;119(2): e348-59.

Bahar A, Abusham A, Eskandar M, Sobande A, Alsunaidi M. Risk factors and pregnancy outcome in different types of placenta previa. *J Obstet Gynaecol Can.* 2009;31(2):126-31.

Bailey RE. Intrapartum Fetal Surveillance. ALSO Guidelines. Available at:

http://www.aafp.org/online/etc/medialib/aafp_org/documents/cme/courses/clin/also/fetalsurveillance.Par.0001.File.tmp/FetalSurveillance20070830final.pdf

Balsells M, Garcia-Patterson A, Gich I, Corcoy R. Maternal and fetal outcome in women with type 2 versus type 1 diabetes mellitus: a systematic review and metaanalysis. *J Clin Endocrinol Metab.* 2009;94(11):4284-91.

Barbosa-Cesnik C, Schwartz K, Foxman B. Lactation mastitis. *JAMA.* 2003;289(13):1609-1612.

Bateman BT, Berman MF, Riley LE, Leffert LR. The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesth Analg.* 2010;110(5):1368-73.

Beeckman K, van De Putte S, Putman K, Louckx F. Predictive social factors in relation to preterm birth in a metropolitan region. *Acta Obstet Gynecol Scand.* 2009; 88(7):787-92.

Beilin Y, Bodian CA, Weiser J, Hossain S, Arnold I, Feirman DE, Martin G, Holzman I. Effect of labor epidural analgesia with and without fentanyl on infant breast-feeding: A prospective randomized, double-blind study. *Anesthesiology.* 2005; 103:1211-1217.

Ben-David G, Sheiner E; Levy A; Erez O; Mazor M. An increased risk for non allo-immunization related intrauterine fetal death in RhD-negative patients. *J matern Fetal Neonatal Med.* 2008; 21(4):255-9.

Ben-Haroush A, Yogeve Y, Bar J, Glickman H, Kaplan B, Hod M. Indicated labor induction with vaginal prostaglandin E2 increases the risk of cesarean section even in multiparous women with no previous cesarean section. *J Perinat Med.* 2004;32(1):31-6.

Berg CJ, Chang J, Callaghan WM, Whitehead SJ. Pregnancy-related mortality in the United States, 1991-1997. *Obstet Gynecol.* 2003;101(2):289-96.

Bhat RY, Rao A. Meconium-stained amniotic fluid and meconium aspiration syndrome: A prospective study. *Annals of Tropical Paediatrics.* 2008; 28:199-203.

Bjørstad AR, Irgens-Hansen K, Daltveit AK, Irgens LM. Macrosomia: mode of delivery and pregnancy outcome. *Acta Obstet Gynecol Scand.* 2010;89(5):664-9.

Black M, Shetty A, Bhattacharya S. Obstetric outcomes subsequent to intrauterine death in the first pregnancy. *BJOG.* 2008;115:269-274.

Blume SL, Casey BM, Schaffer JJ, McIntire DD, Leveno KJ. A randomized trial of coached versus uncoached maternal pushing during the second stage of labor. *AJOG;* 2005; 194:10-13.

Blume HK, Li CI, Loch CM, Koepsell TD. Intrapartum fever and chorioamnionitis as risks for encephalopathy in term newborns: a case-control study. *Developmental Medicine and Child Neurology*. 2008;50(1):19-24.

Bodner K, Wierrani F, Grünberger W, Bodner-Adler B. Influence of the mode of delivery on maternal and neonatal outcomes: a comparison between elective cesarean section and planned vaginal delivery in a low-risk obstetric population. *Archives of Gynecology Obstetrics*. 2011;Jun;283(6):1193-8. Epub 2010 May 27.

Botet F, Figueras J, Carbonnell-Estrany X, Arca G. Effect of maternal clinical chorioamnionitis on neonatal morbidity in very-low birthweight infants: a case – control study. *J Perinatal Medicine*. 2010;38(3):269-73.

Boy A, Salihi HM. Intimate partner violence and birth outcomes: a systematic review. *Int J Fertil Womens Med*. 2004;49(4):159-64.

Boyd HA, Poulson G, Wohlfahrt J, Murray JC, Feenstra B, Melbye M. Maternal contributions to preterm delivery. *Am J Epidemiol*. 2009; 170(11):1358-64.

Brabin BJ, Hakimi M, Pelletier D. An analysis of anemia and pregnancy-related maternal mortality. *J Nutr*. 2001;131(2S-2):604S-614S.

Brady MT. Health care-associated infections in the neonatal intensive care unit. *Am J Infect Control*. 2005;33(5):268-75.

Briggs G, Freeman RK, Yaffe SJ. *Drugs in pregnancy & lactation: A reference guide to fetal and neonatal risk*. 8th Ed. Williams & Wilkins. 2008

Brown M, Mackenzie C, Dunsmuir W, Roberts L, Iken K, Matthews J, Mangos G, Davis D. Can we predict recurrence of preeclampsia or gestational hypertension? *BJOG* 2007;7114:984-993.

Bryant AS, Worjolah A, Caughen AB, Washington AE. Racial/ethnic disparities in obstetric outcomes and care: prevalence and determinants. *Am J Obstet Gynecol*. 2010; 202(4):335-43.

Bussel JB. Fetal and neonatal cytopenias: what have we learned? *Am J Perinatol*. 2003;20(8):425-31.

Burd L, Roberts D, Olson M, Odendaal H. Ethanol and the placenta: A review. *J Matern Fetal Neonatal Med*. 2007;29(5):361-75.

Cammu H, Martens G, Ruysinck G, Amy JJ. Outcome after elective labor induction in nulliparous women: matched cohort study. *Am J Obstet Gynecol*. 2002; 186: 240-4.

Campbell D, Scott KD, Klaus MH, Falk M. Female relatives or friends trained as labor doulas” Outcomes at 6 to 8 weeks postpartum. *Birth*. 2007; 34(3):220-227.

Carfoot S, Williamson P, Dickson R. A systematic review of randomized controlled trials evaluating the effect of mother/baby skin-to-skin care on successful breastfeeding. *Midwifery*. 2003;19:148-55.

Carroli G, Mignini L. Episiotomy for vaginal birth. *The Cochrane Database of Systematic Reviews*, 2009, Issue 1, Art NO: CD000081. DOI: 10.1002/14651858.CD000081.pub2.

Carroli G, Rooney C, Villar J. How effective is antenatal care in preventing maternal mortality and serious morbidity? An overview of the evidence. *Paediatr Perinat Epidemiol*. 2001;15 Suppl 1:1-42.

Caughey AB, Sundaram V, Kaimal AJ, Gienger A, Cheng YW, McDonald KM, Shaffer BL, Owens DK, Bravata DM. Systematic review: Elective induction of labor versus expectant management of pregnancy. *Ann Intern Med*. 2009; 151:252-263.

Cecatti JG, Correa-Silva EP, Milanez H, Morais SS, Souza JP. The associations between inter-pregnancy interval and maternal and neonatal outcomes in Brazil. *Matern Child Health J*. 2008; 12(2):275-81.

Cederholm M, Haglund B, Axelsson O. Maternal complications following amniocentesis and chorionic villus sampling for prenatal karyotyping. *BJOG*. 2003;110(4):392-9.

Cederhold M, Haglund B, Axelsson O. Infant morbidity following amniocentesis and chorionic villus sampling for prenatal karyotyping. *BJOG*. 2005;112(4):394-402.

Cedergren MI. Non-elective caesarean delivery due to ineffective uterine contractility or due to obstructed labour in relation to maternal body mass index. *Eur J Obstet Gynecol Reprod Biol*. 2009; 145(2):163-6.

Chaim W, Bashiri A, Bar-David J, Shoham-Vardi I, Mazor M. Prevalence and clinical significance of postpartum endometritis and wound infection. *Infect Dis Obstet Gynecol*. 2000;8:77-82.

Chalupczak P, Kolasinska-Kloch W, Jach R, Basta A. Pregnancy in patients with heart disease. *Clin Exp Obstet Gynecol*. 2004;31(4):271-3.

Chappell LC, Enye S, Seed P, Briley AL, Poston L, Shennan AH. Adverse perinatal outcomes and risk factors for preeclampsia in women with chronic hypertension: a prospective study. *Hypertension*. 2008; 51(4):1002-9.

Charpak N, Ruiz-Pelaez JG, Figueroa de CZ, Charpak Y. A randomized, controlled trial of kangaroo mother care. Results of follow-up at 1 year of corrected age. *Pediatrics*. 2001;108(5):1072-79.

Christou H, Brodsky D. Lung injury and bronchopulmonary dysplasia in newborn infants. *J Intensive Care Med*. 2005;20(2):76-87.

- Chu SY et al. Gestational weight gain by body mass index among US women delivering live infants, 2004-2005: fueling future obesity. *Am J Obstet Gynecol*. 2009; 271:271e1-271e7.
- Cleary-Goldman J, Malone FD, Vidaver J, Ball RH, Nyberg DA, Comstock CH, Saade GR, Eddleman KA, Klugman S, Dugoff L, Timor-Tritsch IE, Craigo SD, Carr SR, Wolfe HM, Bianchi DW, D'Alton M. Impact of maternal age on obstetric outcome. *Obstet Gynecol*. 2005;105(5 Pt 1):983-90.
- Conde-Agudelo A, Belizan J, Lindmark G. Maternal morbidity and mortality associated with multiple gestations. *Obstet Gynecol*. 2000;95:899-904.
- Conde-Agudelo A, Diaz-Rossello JL, Belizan JM. Kangaroo mother care to reduce mortality in low birthweight infants. In: *The Cochrane Database of Systematic Reviews*, 2003.
- Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *JAMA*. 2006;295(15):1809-23.
- Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: A systematic review. *AJOG*. 2007;196(4):297-308.
- Cornelius MD, Day NL. Developmental consequences of prenatal tobacco exposure. *Curr Opin Neurol*. 2009;22(2):121-5.
- Cotter AM, Martin CM, O'Leary JJ, Daly SF. Increased fetal RhD gene in the maternal circulation in early pregnancy is associated with an increased risk of pre-eclampsia. *BJOG*. 2005;112(5):584-7.
- Crowther CA. Anti-D administration in pregnancy for preventing Rhesus alloimmunisation. In: *The Cochrane Database of Systematic Reviews*, 2003.
- Crowther CA, Hiller JE, Moss JR, McPhee AJ, Jeffries WS, Robinson JS. Effect of treatment of gestational diabetes mellitus on pregnancy outcomes. *N Engl J Med*. 2005;352(24):2477-86.
- Crowther C, Middleton P. Anti-D administration after childbirth for preventing Rhesus alloimmunisation. In: *The Cochrane Database of Systematic Reviews*, 2003.
- Daltveit A, Tollanes MC, Pihlstrom H, Irgens L. Cesarean delivery and subsequent pregnancies. *Obstetrics & Gynecology*. 2008; 111(6):1327-34.
- Dasgupta SJ, Gill AB. Hypotension in the very low birthweight infant: the old, the new, and the uncertain. *Arch Dis Child Fetal Neonatal Ed*. 2003;88(6):F450-4.
- Davies GA, Maxwell C, McLeod L, Gagnon R, Basso M, Bos H, Delisle MF, Farine D, Hudon L, Menticoglou S, Mundle W, Murphy-Kaulbeck L, Ouellet A, Pressey T, Roggensack A, Leduc D, Ballerman C, Biringer A, Duperron L, Jones D, Lee LS, Shepherd D, Wilson K, Society of Obstetricians and Gynaecologists of Canada. Obesity in pregnancy. *J Obstet Gynaecol Can*. 2010;32(3):165-73.

Decca L, Daldoss C, Fratelli N, Lojacono A, Slompo M, Stegher C, Valcamonico A, Frusca T. Labor course and delivery in epidural analgesia: a case-control study. *J Matern Fetal Neonatal Med.* 2004;16(2):115-8.

Declercq E, Barger M, Cabral JH, Evans SR, Kotelchuck M, Simon C, Weiss J, Jeffner LJ. Maternal outcomes associated with planned primary cesarean births compared with planned vaginal births. *Obstet Gynecol.* 2007;109(3):669-677.

De Jonge A, Teunissen TA, LARGRO-JANSSEN AL. Supine position compared to other positions during the second stage of labour : a meta-analytic review. *J Psychosomatic Obstetrics and Gynaecology.* 2005;25:35-45.

Delpisheh A, Brabin L, Attia E, Brabin BJ. Pregnancy late in life: a hospital-based study of birth outcomes. *J Women's Health.* 2008; 17(6):965-70.

Delpisheh A, Brabin L, Drummond S, Brabin BJ. Prenatal smoking exposure and asymmetric fetal growth restriction. *Ann Hum Biol.* 2008;35(6):573-83.

Demissie K, Rhoads GG, Smulian JC, Balasubramanian BA, Gandhi K, Joseph KS, Kramer M. Operative vaginal delivery and neonatal and infant adverse outcomes: population based retrospective analysis. *BMJ.* 2004;329(7456):24-9.

Dodd JM, Crowther CA, Huertas E, Guise JM, Horey D. Planned elective repeat caesarean section versus planned vaginal birth for women with a previous caesarean birth. *Cochrane Database Syst Rev.* 2004, Issue 4. Art. NO: CD004224. DOI: 10.1002/14651858.CD004224.pub.2

Donnelly A, Snowden HM, Renfrew MJ, Woolridge MW. Commercial hospital discharge packs for breastfeeding women. In: *The Cochrane Database of Systematic Reviews*, 2005.

Dryden C, Young D, Hepburn M, Mactier H. Maternal methadone use in pregnancy: factors associated with the development of neonatal abstinence syndrome and implications for healthcare resources. *BJOG.* 2009;116(5):665-71.

Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol.* 2009; 33(3):130-7.

Dyson L, McCormick F, Renfrew MJ. Interventions for promoting the initiation of breastfeeding. In: *The Cochrane Database of Systematic Reviews*, 2005.

Eddleman KA., Malone FD, Sullivan L, Dukes K, Berkowitz RL, Kharbutli Y et al., Pregnancy loss rates after midtrimester amniocentesis. *Obstet Gynecol.* 2006;108(5):1067-72.

Elbourne D, Wiseman RA. Types of intra-muscular opioids for maternal pain relief in labour. In: *The Cochrane Database of Systematic Reviews*, 2003

Elsedeek M, Awad E, Elsebaey S. Evaluation of postpartum blood loss after misoprostol-induced labour. *BJOG.* 2009; 116:431-435.

El Marroun H, Tiemeier H, Steegers EA, Jaddoe VW, Hofman A, Verhulst A, Verhulst FC, van den Brink W, Huizink AC. Intrauterine cannabis exposure affects fetal growth trajectories: The Generation R Study. *J Am Acad Child Adolesc Psychiatry*. 2009; 48(12):1173- 1181.

Esplin MS, O'Brien E, Fraser A, Kerber RA, Clark E, Simonsen SE, Holmgren C, Mineau GP, Varner MW. Estimating recurrence of spontaneous preterm delivery. *Obstet Gynecol*. 2008; 112(3):516-23.

Erez O, Shoham-Vardi I, Sheiner E, Dukler D, Bashiri A, Mazor M. Hydramnios and small for gestational age are independent risk factors for neonatal mortality and maternal morbidity. *Arch Gynecol Obstet*. 2005; 271:296-301.

Esakoff TF, Cheng YW, Sparks TN, Caughey AB. The association between birthweight 4000g or greater and perinatal outcomes in patients with and without gestational diabetes mellitus. *Am J Obstet Gynecol*. 2009;200(6):672.31-4.

Evron S, Dimitrochenko V, Khazin V, Sherman A, Sadan O, Boaz M, Ezri T. The effect of intermittent versus continuous bladder catheterization on labor duration and postpartum urinary retention and infection: a randomized trial. *J Clin Anesth*. 2008;20(8):567-72.

Ezra Y, Michaelson-Cohen R, Abramov Y, Rojansky N. Prelabor rupture of the membranes at term: when to induce labor? *Eur J Obstet Gynecol Reprod Biol*. 2004;115(1):23-7.

Fernando RJ, Sultan AHH, Kettle C, Thakar R, Radley S. Methods of repair for obstetric and spinchter injury. *Cochrane Database Syst Rev*. 2006; Issue 2, Art No: CD002866. DOI: 10.1002/14651858

Fischer MJ, Lehnerz SD, Hebert JR, Parikh CR. Kidney disease is an independent risk factor for adverse fetal and maternal outcomes in pregnancy. *Am J Kidney Dis*. 2004;43(4):415-23.

Frederick IO, Williams MA, Sales AE, Martin DP, Killien M. Pre-pregnancy body mass index, gestational weight gain, and other maternal characteristics in relation to infant birth weight. *Matern Child Health J*. 2008; 12(5):557-67.

Fleming VE, Hagen S, Niven C. Does perineal suturing make a difference? The SUNS trial. *Br J Obstet Gynaecol*. 2003;119(7):684-9

Flenady V, King J. Antibiotics for prelabor rupture of membranes at or near term. In: *The Cochrane Database of Systematic Reviews*, 2003

Floyd RL, Weber MK, Denny C, O'Connor MJ. Prevention of fetal alcohol spectrum disorders. *Dev Disabil Res Rev*. 2009;15(3):193-9.

Forna F, Jamieson DJ, Sanders D, Lindsay MK. Pregnancy outcomes in foreign-born and US-born women. *Int J Gynaecol Obstet*. 2003;83(3):257-65.

Forrester MB, Merz RD. risk of selected birth defects with prenatal illicit drug use, Hawaii, 1986-2002. *J Toxicol Environ Health A*. 2007;70(1):7-18.

Foxman B. Epidemiology of urinary tract infections: incidence, morbidity and economic costs. *Am J Med.* 2002;113 Suppl 1A:5S-13S.

French L. Prevention and treatment of postpartum endometritis. *Curr Women's Health Rep.* 2003;3(4):274-9.

French LM, Smaill FM. Antibiotic regimens for endometritis after delivery. In: *The Cochrane Database of Systematic Reviews*, 2002.

Froen JF, Gardosi JO, Thurmann A, Francis A, Stray-Pedersen B. Restricted fetal growth in sudden intrauterine unexplained death. *Acta Obstet Gynecol Scand.* 2004;83(9):801-7.

Galyean AM, Dagrew DC, Bush MC, Kurtzman JT. Previous cesarean section and the risk of postpartum maternal complications and adverse neonatal outcomes in future pregnancies. *J Perinatol.* 2009; 29:726-730.

Gaugler-Senden IP, Berends AL, de Groot CJ, Steegers EA. Severe, very early onset preeclampsia: subsequent pregnancies and future parental cardiovascular health. *Eur J Obstet Gynecol Reprod Biol.* 2008;140(2):171-7 Epub2008 Jun 20.

Getahun D, Amre D, Rhoads GG, Demissie K. Maternal and obstetric risk factors for sudden infant death syndrome in the United States. *Obstet Gynecol.* 2004;103(4):646-52.

Getahun D, Ananth CV, Kinzler WL. Risk factors for antepartum and intrapartum stillbirth: a population based study. *Am J Obstet Gynecol.* 2007;196(6):499-507.

Getahun D, Oyelese Y, Salihu HM, Ananth CV. Previous cesarean delivery and risks of placenta previa and placental abruption. *Obstet Gynecol.* 2006; 107(4):771-8.

Gilbert WM, Young AL, Danielsen B. Pregnancy outcomes in women with chronic hypertension: a population-based study. *J Reprod Med.* 2007; 52(11):1046-51.

Glantz JC. Term labor induction compared with expectant management. *Obstet Gynecol.* 2010; 115:70-76.

Goedhart G, van Eijsden M, van der Wal, MF, Bonsel GJ. Ethnic differences in preterm birth and its subtypes: the effect of a cumulative risk profile. *BJOG.* 2008;115(6):710-9.

Golden S, Bennett W, Baptist-Roberts K, Wilson L, Barone B, Gary T, Bass E, Nicholson W. Antepartum glucose tolerance test results as predictors of Type 2 diabetes mellitus in women with a history of gestational diabetes mellitus: A systematic review. *Gender Medicine.* 2009;6:109-122.

Goldenberg RL, Culhane JF, Iams JD, Romero R, Epidemiology and causes of preterm birth. *Lancet* 2008; 371(9696):75-84.

Gourounti K, Sandall J. Admission cardiotocography versus intermittent auscultation of fetal heart rate: effects on neonatal Apgar score, on the rate of caesarean sections and on the rate of instrumental delivery – a systematic review. *Int J Nursing Stud.* 2007;44(6):1029-35.

Graneli SL, McGrath JM. Neonatal seizures: diagnosis, pharmacologic interventions, and outcomes. *J Perinat Neonatal Nurs.* 2004;18(3):275-87.

Gray R, Quigley MA, Hockley C, Kurinczuk JJ, Goldacre M, Brocklehurst P. Cesarean delivery and risk of stillbirth in subsequent pregnancy: a retrospective cohort study in an English population. *BJOG.* 2007;114(3):264-70.

Grivell RM, Wong L, Bhatia V. Regimens of fetal surveillance for impaired fetal growth. *Cochrane Database of Systematic Reviews.* 2009. Issue 1. Art. No.: CD007113. DOI: 10.1002/14651858.CD007113.pub2.

Grobman W, Lai Y, Landon MB., et al. Can a prediction model for vaginal birth after cesarean also predict the probability of morbidity related to a trial of labor ? *Am J Obstet Gynecol.* 2009;200:56.e1-56.e6

Grobman WA, Auger M, Shulman LP, Elias S. The association between chorionic villus sampling and preeclampsia. *Prenatal Diagnosis.* 2009; 29(8):800-3.

Grobman WA, Gersnoviez R, Landon MB, Spong CY, Leveno KJ, Rouse DJ, Varner MW, Moawad AH, Caritis SN, harper M, Wapner RJ, Sorokin Y, Miodovnik M, Carpenter M, O'Sullivan MJ, Sibai BM, Langer O, Thorp JM, Ramin SM, Mercer BM. Pregnancy outcomes for women with placenta previa in relation to the number of prior cesarean deliveries. *Obstet Gynecol.* 2007;110(6):1249-55.

Gul A, Cebeci A, Aslan H, Polat I, Ozdemir A, Ceylan Y. Perinatal outcomes in severe preeclampsia-eclampsia with and without HELLP syndrome. *Gynecol Obstet Invest.* 2005;59(2):113-8.

Guner YS, Griedlich P, Wee CP, Dorey F, Camerini V, Upperman JS. State-based analysis of necrotizing enterocolitis outcomes. *J Surgical Research.* 2009; 157(1):21-9.

Gupta JK, Hofmeyr GJ, Smyth RMD. Position in the second stage of labour for women without epidural anaesthesia. *Cochrane Database of Systematic Reviews.* 2004. Issue 1. Art. No.:CD002006. DOI: 10.1002/14651858.CD002006.pub.2.

Haeri S, Shauer M, Dale M, Leslie J, Baker AM, Saddlemire S, Boggess K. Obstetric and newborn infant outcomes in human immunodeficiency virus-infected women who receive highly active antiretroviral therapy. *Am J Obstet Gynecol.* 2009; 201(3):315.e1-5.

Hansen A, Wisborg K, Uldbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *British Medical Journal.* 2008; 336:85-87.

- Hanson S, Clark S, Foster J. Active pushing vs passive fetal descent in the second state of labor: A randomized controlled trial. *Obstet Gynecol.* 2002;99:29-34.
- Harlev A, Levy A, Zaulan Y, Koifman A, Mazor M, Wiznitzer A, Faizayev E, Sheiner A. Idiopathic bleeding during the second half of pregnancy as a risk factor for adverse perinatal outcome. *J Matern Fetal Neonatal Med.* 2008;21(5):331-5.
- Harlow BL, Vitonis AF, Sparen P, Cnattingius S, Joffe H, Hultman CM. Incidence of hospitalization for postpartum psychotic and bipolar episodes in women with and without prior pre-pregnancy or prenatal psychiatric hospitalizations. *Arch Gen Psychiatry.* 2007;64(1):42-8.
- Harrington DJ, Redman CW, Moulden M, Greenwood CE. Long-term outcome in surviving infants with Apgar zero at 10 minutes: a systematic review of the literature and hospital-based cohort. *Am J Obstet Gynecol.* 2007;195(5):463-4.
- Hartmann K, Viswanathan M, Palmieri R, Gertlehner G, Thorp J, Lohr KN. Outcomes of routine episiotomy: a systematic review. *JAMA.* 2005;293:2141-8.
- Havens JR, Simmons LA, Shannong LM, Hansen WF. Factors associated with substance use during pregnancy: results from a national sample. *Drug Alcohol Depend.* 2009; Jan 1;99(1-3):89-95.
- Haydon ML, Gorenberg DM, Nageotte MP, Ghamsary M, Rumney PJ, Patillo C, Garite TJ. The effect of maternal oxygen administration on fetal pulse oximetry during labor in fetuses with nonreassuring fetal heart rate patterns. *Am J Obstet Gynecol.* 2006; 195:735-8.
- Heinemann J, Gillen G, Sanchez-Ramos L, Kaunitz AM. Do mechanical methods of cervical ripening increase infectious morbidity? A systematic review. *Am J Obstet Gynecol.* 2008; 199(2):177-188.
- Henriksen TB, Hjollund NH, Jensen TK, Bonde JP, Andersson AM, Kolstad H, Ernst E, Giwercman A, Skakkebaek NE, Olsen J. Alcohol consumption at the time of conception and spontaneous abortion. *Am J Epidemiol.* 2004;160(7):661-7.
- Herbst MA, Mercer BM, Beazley D, Meyer N, Carr T. Relationship of prenatal care and perinatal morbidity in low-birth-weight infants. *Am J Obstet Gynecol.* 2003;189(4):930-3.
- Hill JB, Alexander JM, Sharma SK, McIntire DD, Leveno KJ. A comparison of the effects of epidural and meperidine analgesia during labor on fetal heart rate. *Obstet Gynecol.* 2003;102(2):333-7.
- Hill JB, Sheffield JS, McIntire DD, Wendel GD Jr. Acute pyelonephritis in pregnancy. *Obstet Gynecol.* 2005;105(1):18-23.
- Hizkiyahu R, Levy A, Sheiner E. Pregnancy outcome of patients with schizophrenia. *Am J Perinatol.* 2010;27(1):19-23.
- Hoch M, Netz H. Heart failure in pediatric patients. *Thorac Cardiovasc Surg.* 2005;53 Suppl 2:S129-34.

- Hodnett ED. Continuity of caregivers for care during pregnancy and childbirth In: The Cochrane Database of Systematic Reviews, 2003.
- Hodnett ED, Gates S, Hofmeyr GJ, Sakala A. Continuous support for women during childbirth. The Cochrane Database of Systematic Reviews. 2007; Issue 3.
- Hoffman JA, Mason EO, Schutze GE, Tan TQ, Barson WJ, Givner LB, Wald ER, Bradley JS, Yogeve R, Kaplan SL. Streptococcus pneumoniae infections in the neonate. Pediatrics. 2003;112(5):1095-102.
- Holmes H, Lo J, McIntire D, Casey B. Prediction of diabetes recurrence in women with class A (diet treated) gestational diabetes. Am J Perinatol. 2010;27(1):47-52.
- Holzman C, Lin X, Senagore P, Chung H. Histologic chorioamnionitis and preterm delivery. Am J Epidemiol. 2007;166(7):786-94.
- Hohmann-Marriott B. The couple context of pregnancy and its effects on prenatal care and birth outcomes. Matern child Health J. 2009; 13(6):745-54.
- Horvath T, Madi B, Iuppa IM, Kennedy GE, Rutherford G, Read JS. Interventions for preventing late postnatal mother-to child transmission of HIV. Cochrane Database Syst Rev. 2009 Jan. 21(1):CD006734.
- Howard LM, Goss C, Leese M, Thornicroft G. Medical outcome of pregnancy in women with psychotic disorders and their infants in the first year after birth. Br J Psychiatry. 2003;182:63-7.
- Howell CJ. Epidural versus non-epidural analgesia for pain relief in labour. In: The Cochrane Database of Systematic Reviews, 2003.
- Howell CJ, Dean T, Lucking L, Dziedzic K, Jones, PW, Johanson RB. Randomised study of long term outcome after epidural versus non-epidural analgesia during labor. BMJ. 2002; 325(7360):357.
- Imbasciati E, Tincani A, Gregorini G, Doria A, Moroni G, Cabiddu G, Marcelli D. Pregnancy in women with pre-existing lupus nephritis: predictors of fetal and maternal outcome. Nephrol Dial Transplant. 2009;24(2):519-25.
- Jablensky AV, Morgan V, Zubrick SR, Bower C, Yellachich LA. Pregnancy, delivery, and neonatal complications in a population cohort of women with schizophrenia and major affective disorders. Am J Psychiatry. 2005;162(1):79-91.
- Jaddoe VW, Bakker R, Hofman A, Mackenbach JP, Moll HA, Steegers EA, Witteman JC. Moderate alcohol consumption during pregnancy and the risk of low birth weight and preterm birth. The generation R study. Ann Epidemiol. 2007;17(10):834-40.
- Jain S, Arya VK, Gopalan S, Jain V. Analgesic efficacy of intramuscular opioids versus epidural analgesia in labor. Int J Gynaecol Obstet. 2003;83(1):19-27.

Jansen AJ, van Rhenen DJ, Steegers EA, Duvekot JJ. Postpartum hemorrhage and transfusion of blood and blood components. *Obstet Gynecol Surv.* 2005;60(10):663-71.

Johansson AL, Dickman PW, Kramer MS, Cnattingius S. Maternal smoking and infant mortality: does quitting smoking reduce the risk of infant death? *Epidemiology.* 2009;20(4):590-7.

Johnson KJ, Soler JT, Puumala SE, Ross JA, Spector LG. Parental and infant characteristics and childhood leukemia in Minnesota. *BMC Pediatr.* 2008; 25:8:7

Kahana B, Sheiner E, Levy A, Lazer S, Mazor M. Umbilical cord prolapse and perinatal outcome. *Int J Gynaecol Obstet.* 2004;84(2):127-32.

Kassanos D, Siristatidis C, Vitoratos N, Salamalekis E, Creatas G. The clinical significance of Doppler findings in fetal middle cerebral artery during labor. *Eur J Obstet Gynecol Reprod Biol.* 2003;109(1):45-50.

Kaupila A, Kyyrönen P, Hinkula M, Pukkala E. Birth intervals and breast cancer risk. *Br. J Cancer.* 2009; 101(7):1213-7.

Kaye DK, Mirembe FM, Bantebya G, Johannsson A, Eksrom AM. Domestic violence during pregnancy and risk of low birthweight and maternal complications: a prospective cohort study at Mulago Hospital, Uganda. *Trop Med Int Health.* 2006;11(10):1576-84.

Kennare R, Rucker G, Heard A, Chan A. Risks of adverse outcomes in the next birth after a first cesarean delivery. *Obstet Gynecol.* 2007;109 (2 Part 1):270-6

Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PFA. WHO analysis of causes of maternal death: a systematic review. *Lancet.* 2006;367:1066-1074.

Khashu M, Narayanan M, Bhargava S, Osioviich H. Perinatal outcomes associated with preterm birth at 33 to 35 weeks' gestation: a population-based cohort study. *Pediatrics.* 2009;123(1):109-13.

Kidanto HL. Risks for preterm delivery and low birth weight are independently increased by severity of maternal anaemia. *S Afr Med J.* 2009; 99(2):98-102.

Kiely M, El-Mohandes AA, El-Khorazaty MN, Gantz MG. An integrated intervention to reduce intimate partner violence in pregnancy: a randomized controlled trial. *Obstet Gynecol.* 2010;115(2 Part 1):273-83.

Kiernan K, Pickett KE. Marital status disparities in maternal smoking during pregnancy, breastfeeding and maternal depression. *Soc Sci med.* 2006; 63(2):335-46.

Kitzmilller JL, Block JM, Brown FM, Catalano PM, Conway DL, Coustan DR et al., Managing preexisting diabetes for pregnancy. Summary of evidence and consensus recommendations for care. *Diabetes Care.* 2008;31(5):1060-79.

Knight M. Eclampsia in the United Kingdom 2005. *BJOG*. 2007; 114(9):1072-8.

Knight M, Kurinczuk JJ, Spark P, Brocklehurst P, United Kingdom Obstetric Surveillance System Steering Committee. Cesarean delivery and peripartum hysterectomy. *Obstet Gynecol*. 2008;111(1):97-105.

Koelewijn JM, de Haas M, Vrijkotte TG, van der Schoot CE, Bonsel FH. Risk factors for RhD immunization despite antenatal and postnatal anti-D prophylaxis. *BJOG*. 2009;116(10):1307-14. Epub 2009 Jun 17.

Konduri GG. New approaches for persistent pulmonary hypertension of newborn. *Clin Perinatol*. 2004;31(3):591-611.

Koifman A, Levy A, Zaulan Y, harlev A, Mazor M, Wiznitzer A, Sheiner E. The clinical significance of bleeding during the second trimester of pregnancy . *Arch Gynecol Obstet*. 2008; 278(1):47-51.

Kovac M, Mitic G, Mikovic Z, Djordjevic V, Savic O, Mandic V, Rakicevic L, Antonijevic N, Radojkovic D. Thrombophilia in women with pregnancy-associated complications: Fetal loss and pregnancy-related venous thromboembolism. *Gynecol Obstet Invest*. 2010;69:233-238.

Kristensen J, Vestergaard M, Wisborg K, Kesmodel U, Secher NJ. Pre-pregnancy weight and the risk of stillbirth and neonatal death. *BJOG*. 2005;112(4):403-8.

Kwak S, Kim H, Choi S, Lim S, Cho Y, Park K, Jang H, Kim M, Cho N, Metzger B. Subsequent pregnancy after gestational diabetes mellitus: frequency and risk factors for recurrence in Korean women. *Diabetes Care*. 2008;31(9):1867-1871.

Lai , CY, Cheung HW, Hsi Lao TT, Lau TK, Leung TY. Is the policy of restrictive episiotomy generalisable? A prospective observational study. *J Maternal-Fetal & Neonatal Med*. 2009;22(12):1116-21.

Langer O, Yogev Y, Most O, Xenakis EM. Gestational diabetes: the consequences of not treating. *Am J Obstet Gynecol*. 2005;192(4):989-97.

Lapolla A, Dalfra MG, DiCianni G, Bonomo M, Parretti E, Mello G. A multicenter Italian study on pregnancy outcome in women with diabetes. *Nutr Metab Cardiovasc Dis*. 2008;18(4):291-7.

Larma AJ, Silva A, Holcroft C, Thompson R, Donohue P, Graham E. Intrapartum electronic fetal heart rate monitoring and the identification of metabolic acidosis and hypoxic-ischemic encephalopathy. *Am J Obstet Gynecol*. 2007;197:301e1-e8.

Lawrence A, Lewis L, Hofmeyr GH, Dowswell R, Styles C. Maternal positions and mobility during first stage labour. *Cochrane Database of Systematic Reviews*. 2009. Apr 15; (2); Cochrane AN: CD003934;

Lee C, Lin HC. Maternal bipolar disorder increased low birthweight and preterm births: a nationwide population-based study. *J Affect Disord.* 2010;121(1-2):100-5.

Lin HC, Chen IJ, Chen YH, Lee HC, Wu FJ. Maternal schizophrenia and pregnancy outcome: does the use of antipsychotics make a difference? *Schizophr Res.* 2010;116(1):55-60.

Lin C, Raynor BD. Risk of uterine rupture in labor induction of patients with prior cesarean section: an inner city hospital experience. *Am J Obstet Gynecol.* 2004;190(5):1476-8.

Liu BY, Wang CC, Lau TK, Chu CY, Phil M, Pang CP, Rogers MS, Leung TN. Meconium-stained liquor during labor is associated with raised neonatal cord blood 8-iso-prostaglandin F₂alpha concentration. *Am J Obstet Gynecol.* 2005;192(1):289-94.

Lobato G, Soncini CS. Relationship between obstetric history and Rh(D) alloimmunization severity. *Arch gynecol Obstet.* 2008; 277(3):245-8. Epub 2007 Aug 31.

Locatelli A, Regalia AL, Patregnani C, Ratti M, Toso L, Ghidini A. Prognostic value of change in amniotic fluid color during labor. *Fetal Diagn Ther.* 2005;20(1):5-9.

Lone FW, Qureshi RN, Emanuel F. Maternal anaemia and its impact on perinatal outcome. *Trop Med Int Health.* 2004; 9(4):486-90.

Lone F, Sultan AH, Thakar R, Beggs A. risk factors and management patterns for emergency obstetric hysterectomy over 2 decades. *Int J Gynaecol Obstet.* 2010;109(1):12-5. Epub 2009 Dec 1.

Luders C, Martins Castro MC, Tital SM, De Castro I, Elias RM, Abensur H, Romão JE Jr. Obstetric outcome in pregnant women on long-term dialysis: A case series. *Am J Kidney Dis.* 2010. Article 10 (in press)

Lundquist M, Olsson A, Nissen E, Norman M. Is it necessary to suture all lacerations after a vaginal delivery? *Birth.* 2000;27(2):79-85.

Lutgendorf MA, Busch JM, Doherty DA, Conza LA, Moone SO, Magann EF. Prevalence of domestic violence in a pregnant military population. *Obstet Gynecol.* 2009; 113(4):866-72.

Luo ZC, Wilkins R, Kramer MS. Disparities in pregnancy outcomes according to marital and cohabitation status. *Obstet Gynecol.* 2004;103(6):1300-7.

Lykke JA, Langhoff-Roos J, Lockwood CJ, Triche EW, Paidas MJ. Mortality of mothers from cardiovascular and non-cardiovascular causes following pregnancy complications in first delivery. *Paediatric and Perinatal Epidemiol* May. 2010 DOI: 10.1111/j.1365-3016.2010.01120.x

Lykke JA, Langhoff-Roos J, Sibai BM, Funai EF, Triche EW, Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and Type 2 diabetes mellitus in the mother. *Hypertension*. 2009; 53(6):944-51.

Lykke J, Paidas M, Langhoff-Roos J. Recurring complications in second pregnancy. *Obstet Gynecol*. 2009; 113(6):1217-1224

MacCabe JH, Martinsson L, Lichtenstein P, Nilsson E, Cnattingius S, Murray RM, Hultman CM. Adverse pregnancy outcomes in mothers with affective psychosis. *Bipolar Disorders*. 2007;9(3):305-309.

MacDorman MF, Mathews TJ. The challenge of infant mortality: have we reached a plateau? *Public Health Rep*. 2009;124(5):670-81.

Macones GA, Cahill A, Pare E, Stamilio DM, Ratcliffe S, Stevens E, Sammel M, Peipert J. Obstetric outcomes in women with two prior cesarean deliveries: is vaginal birth after cesarean delivery a viable option? *Am J Obstet Gynecol*. 2005;192(4):1223-8.

Maconochie N, Doyle P, Prior S, Simmons R. Risk factors for first trimester miscarriage – results from a UK-population-based case-control study. *BJOG*. 2007;114(2):170-86.

Madan I, Romero R, Kusanovic JP, Mittal P, Chaiworapongsa T, Dong Z, Mazaki-Tovi S, Vaisbuch E, Alpay Savasan Z, Yeo L, Kim CJ, Hassan SS. The frequency and clinical significance of intra-amniotic infection and/or inflammation in women with placenta previa and vaginal bleeding: an unexpected observation. *J Perinatal Med*. 2010;38(3):275-9.

Magann EP, Channan SP, Bofill JA, Waddell D, Rust OA, Morrison JC. Maternal morbidity and mortality associated with intrauterine fetal demise: five-year experience in a tertiary referral hospital. *South Med J*. 2001;94(5):493-5.

Malik S, Cleves MA, Honein MA, Romitti PA, Botto LD, Yang S, Hobbs CA; National Birth Defects Prevention Study. Maternal smoking and congenital heart defects. *Pediatrics*. 2008; 121(4):e810-6.

Massad LS, Springer G, Jacobson L, Watts H, Anastos K, Korn A, Cejtin H, Stek A, Young M, Schmidt J, Minkoff H. Pregnancy rates and predictors of conception, miscarriage and abortion in US women with HIV. *AIDS*. 2004;18(2):281-6.

McDonald S, Best C, Lam K. The recurrence risk of severe de novo preeclampsia in singleton pregnancies: a population-based cohort. *BJOG*. 2009;116:1578-1584.

McDonald S, Prendiville WJ, Elbourne D. Prophylactic syntometrine versus oxytocin for delivery of the placenta. In: *The Cochrane Database of Systematic Reviews*, 2002.

McGrath SK, Kennell JH. Continuous labor support for middle-class couples: Effect on cesarean delivery rates. *Birth*. 2008; 35(2):92-97.

McIntyre SH, Newburn-Cook CV, O'Brien B, Demianczuk NN. Effect of older maternal age on the risk of spontaneous preterm labor: a population-based study. *Health Care Women Int.* 2009; 30(8):670-89.

Measey MA, D'Espaignet E, Charles A, Douglass C. Unexplained fetal death: are women with a history of fetal loss at higher risk? *Aus NZ J Obstet Gynaecol.* 2009;49(2):151-7.

Melamed N, Chen R, Soiberman U, Ben-Haroush A, Hod M, Yogev Y. Spontaneous and indicated preterm delivery in pregestational diabetes mellitus: Etiology and risk factors. *Arch Gynecol Obstet.* 2008; 178(2):129-34.

Menezes EV. Reducing stillbirths: prevention and management of medical disorders and infections during pregnancy. *BMC Pregnancy Childbirth.* 2009; 7:9: Suppl 1:S4

Menezes EV, Yakoob MY, Soomro T, Haws RA, Darmstadt GL, Bhutta ZA. Reducing stillbirths: prevention and management of medical disorders and infections during pregnancy. *BMC Pregnancy Childbirth.* 2009; May 7;9 Suppl 1:S4

Metzger BE, Lowe LP, Dyer AR, Trimble ER, Chaovarindr U, Coustan DR et al. Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med* 2008;358:1991-2002.

Moore ER, Anderson GC, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. *The Cochrane Database of Systematic Reviews.* 2007; Issue 4.

Moore LL, Bradless ML, Singer MR, Rothman KJ, Milunsky A. Chromosomal anomalies among the offspring of women with gestational diabetes. *Am J Epidemiol.* 2002;155(8):719-24.

Nabet C, Ancel PY, Burguet A, Kaminski M. Smoking during pregnancy and preterm birth according to obstetric history: French national perinatal surveys. *Paediatr Perinat Epidemiol.* 2005;19(2):88-96.

Natale R, Dodman N. Birth can be a hazardous journey: electronic fetal monitoring does not help. *J Obstet Gynaecol Can.* 2003;25(12):1007-9.

National Collaborating Centre for Women's and Children's Health. Diabetes in pregnancy. Management of diabetes and its complications from pre-conception to the postpartum period. London (UK): National Institute for Health and Clinical Excellence (NICE); 2008 Mar 21, (Clinical guideline, no 63).

Ness RB, Grisso JA, Hirschinger N, Markovic N, Shaw LM, Day NL, Kline J. Cocaine and tobacco use and the risk of spontaneous abortion. *N Engl J Med.* 2002;340(5):333-339.

Normand MC, Damato EG. Postcesarean infection. *J Obstet Gynecol Neonat Nurs.* 2001;30:642-648.

Nystedt A, Edvardsson D, Willman A. Epidural analgesia for pain relief in labour and childbirth – a review with a systematic approach. *J Clin Nurs*. 2004;13(4):455-66.

O’Connell MP, Hussain J, Maclennan FA, Lindow SW. Factors associated with a prolonged second state of labour – a case-controlled study of 364 nulliparous labours. *J Obstet Gynaecol*. 2003;23(3):255-7.

O’Mahony F, Hofmeyr GJ, Menon V. Choice of instruments for assisted vaginal delivery. *Cochrane Database of Systematic Reviews*. 2010, Issue 11. Art. No.:CD005455. DOI:10.1002/14651858.CD005455.pub2.

Odd DE, Rasmussen F, Gunnell D, Lewis G, Whitelaw A. A cohort study of low Apgar scores and cognitive outcomes. *Archives of Disease in Childhood – Fetal & Neonatal Edition*. 2008;93(2): F115-20.

Ogunyemi D, Hernandez-Loera GE. The impact of antenatal cocaine use on maternal characteristics and neonatal outcomes. *J Matern Fetal Neonatal Med*. 2004;15(4):253-9.

Olsen O, Jewell MD. Home versus hospital birth. In: *The Cochrane Database of Systematic Reviews*, 2003.

Onwuhafua PI, Zayyan M, Ozed-Williams I, Ankama M. Fetal outcome following umbilical cord prolapsed in Ahmadu Bello University Teaching Hospital, Kaduna (ABUTHK), northern Nigeria. *Nigerian J Medicine*. 2007;16(4):334-7.

Oron T, Sheiner E, Shoham-Vardi I, Mazor M, Katz M, Hallak M. Risk factors for antepartum fetal death. *J Repro Med*. 2001;46(9):825-30.

Ortibus E. The pregnancy and long-term neurodevelopmental outcome of monochorionic diamniotic twin gestations: a multicenter prospective cohort study from the first trimester onward. *Am J Obstet Gynecol*. 2009; 200(5):494, e1-8.

Osrin D, Vergnano S, Costello A. Serious bacterial infections in newborn infants in developing countries. *Curr Opin Infect Dis*. 2004;17(3):217-24.

Owens R. Intraventricular hemorrhage in the premature neonate. *Neonatal Netw*. 2005;24(3):55-71.

PAHO. Quantifying the benefits of breastfeeding – A summary of the evidence., 2002. Available on-line at: http://www.paho.org/English/HPP/HPN/benefits_of_BF.htm.

Persson M, Norman M, Hanson U. Obstetric and perinatal outcomes in type 1 diabetic pregnancies: A large, population-based study. *Diabetes Care*. 2009;32(11):2005-9.

- Peticca P, Keely EJ, Walker MC, Yang Q, Bottomley J. Pregnancy outcomes in diabetes subtypes: How do they compare? A Province-based study of Ontario, 2005 – 2006. *J Obstet Gynaecol Can.* 2009;31(6):487-496.
- Phipps MG, Blume JD, DeMonner SM. Young maternal age associated with increased risk of postneonatal death. *Obstet Gynecol.* 2002; 100(3):481-6.
- Piccoli GB, Attini R, Vasario E, Conijn A, Biolcati M, D'Amico F, Consiglio V, Bontempo S, Rodros T. Pregnancy and chronic kidney disease: a challenge in all CKD stages. *Clin J Am Soc Nephrol.* 2010; 5(5):844-55.
- Piccoli GB, Conijn A, Consiglio V, Vasario E, Attini R, Deagnostini MC, Bontempo S, Todros T. Pregnancy in dialysis patients: Is the evidence strong enough to lead us to change our counseling policy? *Clin J Am Soc Nephrol.* 2010; 5(1):62-71.
- Pinto SM, Dodd S, Walkiknshaw SA et al., Substance abuse during pregnancy: effect on pregnancy outcomes. *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 2010;150(2):137-41.
- Pitaphrom A, Sukcharoen N. Pregnancy outcome in placental abruption. *J Med Assoc Thailand.* 2006;89(10):1572-8.
- Plunkett J, Borecki I, Morgan T, Stamilio D, Muglia LJ. Population-based estimate of sibling risk for preterm birth, preterm premature rupture of membranes, placental abruption and pre-eclampsia. *BMC Genet.* 20087; Jul 8:9:44.
- Plunkett J, Muglia LJ. Genetic contributions to preterm birth: implications from epidemiological and genetic association studies. *Ann Med.* 2008;40(3):167-95.
- Pulsifer MB, Butz AM, O'Reilly Foran M, Belcher HM. Prenatal drug exposure: effects on cognitive functioning at 5 years of age. *Clin Pediat (Phila).* 2008; 47(1):58-65.
- Rasmussen K. Is there a causal relationship between iron deficiency or iron-deficiency anemia and weight at birth, length of gestation and perinatal mortality? *J Nutr.* 2001;131(2S-2):590S-601S.
- Rasmussen S, Irgens LM, Skjaerven R, Melve KK. Prior adverse pregnancy outcome and the risk of stillbirth. *Obstet Gynecol.* 2009;114(6):1259-70.
- Raatikainen K, Heiskanen N, Heinonen S. Marriage still protects pregnancy. *BJOG.* 2005;112(10):1411-6.
- Raatikainen K, Heiskanen N, Heinonen S. Under-attending free antenatal care is associated with adverse pregnancy outcomes. *BMC Public Health.* 2007; Sept 27;7:268.
- Rahman J, Bhattee G, Rahman MS. Shoulder dystocia in a 16-year experience in a teaching hospital. *J Reprod Med.* 2009; 54(6):378-84.

- Ray JG, Burrows RF, Burrows EA, Vermeulen MJ. MOS HIP: McMaster outcome study of hypertension in pregnancy. *Early Hum Dev.* 2001; 64(2):129-43.
- Revez L. Treatments for iron-deficiency anaemia in pregnancy. *Cochrane Database Syst Rev.* 2007 Apr 18;(2):CD003094.
- Reynolds F, Sharma SK, Seed PT. Analgesia in labour and fetal acid-base balance: a meta-analysis comparing epidural with systemic opioid analgesia. *Br J Obstet Gynaecol.* 2002;109:1344-1353.
- Richardson GA, Goldschmidt L, Larkby C. Effects of prenatal cocaine exposure on growth: a longitudinal analysis. *Pediatrics.* 2007; 120(4): e1017-1027.
- Rich-Edwards JW, Grizzard TA. Psychosocial stress and neuroendocrine mechanisms in preterm delivery. *Am J Obstet Gynecol.* 2005; 192(5 Suppl):S30-5.
- Richter R, Bergmann R, Dudenhausen J. Previous caesarean or vaginal delivery: Which mode is a greater risk of perinatal death at the second delivery? *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 2007;132:51-57.
- Ridgeway JJ, Weyrich DL, Benedetti TJ. Fetal heart rate changes associated with uterine rupture. *Obstet Gynecol.* 2004;103(3):506-12.
- Roberts JE. The “push” for evidence: management of the second stage. *J Midwif Womens Health.* 2002;47(1):2-15.
- Rodriguez A. Maternal pre-pregnancy obesity and risk for inattention and negative emotionality in children. *J Child Psychol Psychiatry.* 2010;51(2):134-43.
- Rogers JM. Tobacco and pregnancy. *Reprod Toxicol.* 2009;28(2):152-60.
- Rojas MA, Kaplan M, Quevedo M, Sherwonit E, Foster LB, Ehrenkranz RA, Mayes L. Somatic growth of preterm infants during skin-to-skin care versus traditional holding: a randomized, controlled trial. *J Dev Behav Pediatr.* 2003;24(3):163-8.
- Ronsmans C, Graham W, The Lancet Maternal Survival Series Steering Group. Maternal mortality: who, when, where, and why. *Lancet.* 2006;368:1189-200.
- Rosen D, Seng J, Tolman R, Mallinger G. Intimate partner violence, depression and posttraumatic stress disorder as additional predictors of low birth weight infants among low-income mothers. *J Interpersonal Violence.* 2007;22:1305-1314.
- Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskiene E, Samuelsson U. Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood. *Diabetes Metab Res Rev.* 2004;20(2):150-7.
- Saleem Z, Rydhström H. Vaginal hematoma during parturition: a population-based study. *Acta Obstet Gynecol Scand.* 2004; 83(6):560-2.

Said JM, Higgins JR, Moses EK, Walker SP, Borg AJ, Monagle PT, Brennecke SP. Inherited thrombophilia polymorphisms and pregnancy outcomes in nulliparous women. *Obstet Gynecol.* 2010; 115(1):5-13.

Salihu HM, Wilson RE, Alio AP, Kirby RS. Advanced maternal age and risk of antepartum and intrapartum stillbirth. *J Obstet Gynaecol Res.* 2008; 34(5):843-50.

Scanlon KS, Yip R, Schieve LA, Cogswell ME. High and low hemoglobin levels during pregnancy: differential risks for preterm birth and small for gestational age. *Obstet Gynecol.* 2000;95 (5 Pt 1):741-8.

Sciscione AC, Landon MB, Leveno KJ, Spong CY, Macpherson C, Varner MW, Rouse DJ, Moawad AH, Caritis SN, Harper M, Sorokin Y, Miodovnik M, Marshall C, Peaceman AM, O'Sullivan MJ, Sibair BM, Langer O, Thorp JM, Ramin SM, Mercer BM. National Institute of Child Health and Human Development (NICHD) Maternal-Fetal Medicine Units Network (MFMU). Previous preterm cesarean delivery and risk of subsequent uterine rupture. *Obstet Gynecol.* 2008; 111(3):648-53.

Schneid-Kofman N, Sheiner E, Levy A. Psychiatric illness and adverse pregnancy outcomes. *Int J Gynaecol Obstet.* 2008;101(1):53-6.

Scholes D, Hooton TM, Roberts PL, Gupta K, Stapleton AE, Stamm WE. Risk factors associated with acute pyelonephritis in healthy women. *Ann Intern Med.* 2005; 142(1):134.

Schonewille H, Klumper FJ, van de Watering LM, Kanhai HH, Brand A. High additional maternal red cell alloimmunization after Thesus- and K-matched intrauterine intravascular transfusions for hemolytic disease of the fetus. *Am J Obstet Gynecol.* 2007; 196(2):143. E1-6.

Schrag S, Schuchat A. Prevention of neonatal sepsis. *Clin Perinatol.* 2005;32(3):601-15.

Sen B, Swaminathan S. Maternal prenatal substance use and behavior problems among children in the U.S. *J Ment Health Policy Econ.* 2007;10(4):189-206.

Schaffer JL, Bloom SL, Casey BM et al., A randomized trial of the effects of coached vs uncoached maternal pushing during the second stage of labor on postpartum pelvic floor structure and function. *AJOG.* 2005; 192(5):1692-1696.

Shand AW, Bell JC, McElduff A, Morris J, Roberts CL. Outcomes of pregnancies in women with pre-gestational diabetes mellitus and gestational diabetes mellitus: a population-based study in New South Wales, Australia, 1998-2002. *Diabetes Medicine.* 2008;25(6):708-15.

Sharma PP, Salihu HM, Ovelese Y, Ananth CV, Kirby RS. Is race a determinant of stillbirth recurrence? *Obstet Gynecol.* 2006;107(2 Pt 1):391-7.

Sharma PP, Shaiju HM, Kirby RS. Stillbirth recurrence in a population of relatively low-risk mothers. *Paediatric and Perinatal Epidemiology.* 2007;21 (Suppl. 1), 24-30.

Sharps P, Laughon K, Giangrande S. Intimate partner violence and the childbearing year: Maternal and infant health consequences. *Trauma, Violence and Abuse*. 2007;6(2):105-116.

Shaw AK, Infante-Rivard C, Morrison HI. Use of medication during pregnancy and risk of childhood leukemia (Canada). *Cancer Causes Control*. 2004;15(9):931-7.

Sheiner E, Shoham-Vardi I, Hallak M, Hershkowitz R, Katz M, Mazor M. Placenta previa: obstetric risk factors and pregnancy outcome. *J Mat Fetal Med*. 2001;10(6):414-9.

Sheiner E, Shoham-Vardi I, Hadar A, Hallak M, Hackmon R, Mazor M. Incidence, obstetric risk factors and pregnancy outcome of preterm placental abruption: a retrospective analysis. *J Mat Fetal Neonat Med*. 2002;11(1):34-9.

Sheiner E, Shohan-Vardi I, Hadar A, Hershkowitz R, Sheiner EK, Mazor M. Accidental out-of-hospital delivery as an independent risk factor for perinatal mortality. *J Reprod Med*. 2002; 47(8):625-30.

Shipp T, Zelop C, Cohen A, Repke J, Lieberman E. Post-cesarean delivery fever and uterine rupture in a subsequent trial of labor. *Obstet Gynecol*. 2003;101(1):136-139.

Sidlik R, Sheiner E, Levy A, Wiznitzer A. Effect of maternal congenital heart defects on labor and delivery outcome: a population-based study. *J Matern Fetal Neonatal Med*. 2007;20(3):211-6.

Silver RM, Landon MB, Rouse DJ, Leveno KJ, Spong CY, Thom EA, et al., Maternal morbidity associated with multiple repeat cesarean deliveries. *Obstet Gynecol*. 2006;107(6):1226-32.

Sikorski J, Renfrew MJ, Pindorial S, Wade A. Support for breastfeeding mothers. In: *The Cochrane Database of Systematic Reviews*, 2005.

Silverman JG, Decker MR, Reed E, Raj A. Intimate partner violence victimization prior to and during pregnancy among women residing in 26 states: associations with maternal and neonatal health. *Am J Obstet Gynecol*. 2006;195(1):140-8.

Simpson KR, James DC. Effects of immediate versus delayed pushing during second-stage labor on fetal well-being: a randomized clinical trial. *Nurs Res*. 2005;54(3):149-57.

Sisco KM, Cahill AG, Stamilio DM, Macones GA. Is continuous monitoring the answer to incidentally observed fetal heart rate decelerations? *J Matern Fetal Neonatal Med*. 2009; 22(5): 405-9.

Siu S, Colman J, Sorensen S, Smallhorn J, Farine D, Amankwah K, Spears J, Sermer M. Adverse neonatal and cardiac outcomes are more common in pregnant women with cardiac disease. *Circulation*. 2002;105:2179-2184.

Siu S, Sermer M, Colman J, Alvarez AN, Mercier L, Morton B, Kells C, et al. Prospective multicenter study of pregnancy outcomes in women with heart disease. *Circulation*. 2001;104:515-521.

Sloan N, Jordan E, Winikoff M. Effects of iron supplementation on maternal hematologic status in pregnancy. *Am J Public Health*. 2002;92(2):268-293.

Smith LM, La Gasse LL, Derauf C, Grant P, Shah R, Arria A, Huestis M, Haning W, Strauss A, Della Grotta S, Liu J, Lester, BM. The infant development, environment, and lifestyle study: effects of prenatal methamphetamine exposure, polydrug exposure, and poverty on intrauterine growth. *Pediatrics*. 2006; 118(3):1149-56.

Smith GC, Shah I, White IR, Pell JP, Dobbie R. Previous preeclampsia, preterm delivery, and delivery of a small for gestational age infant and the risk of unexplained stillbirth in the second pregnancy: a retrospective cohort study, Scotland, 1992-2001. *Am J Epidemiol*. 2007; 165(2):194-202.

Smulian JC, Potash SK, Lai YL, Scorza WE. Appropriateness of antibiotic use in the postpartum period. *J Mat Fetal Med*. 2001;10:312-317.

Smyth RMD, Alldred SK, Markham C. Amniotomy for shortening spontaneous labour. *Cochrane Database of Systematic Reviews*. 2007; Issue 4.

Soll RF. Synthetic surfactant for respiratory distress syndrome in preterm infants. In: *The Cochrane Database of Systematic Reviews*, 2005.

Soong B, Barnes M. Maternal position in midwife-attended birth and perinatal trauma: Is there an association? *Birth*. 2005;32(3):164-169.

Soraisham AS, Singhal N, Mc Millan DD, Sauve RS, Lee SK. A multicenter study on the clinical outcome of chorioamnionitis in preterm infants. *Am J Obstet Gynecol*. 2009; 200(4): 372-e1-6.

Sosa CG, Althabe F, Belizán JM, Buekens P. Risk factors for postpartum hemorrhage in vaginal deliveries in a Latin-American population. *Obstet Gynecol*. 2009; 113(6):1313-9.

Stamilio DM, De Franco E, Paré E, Odibo AO, Peipert JF, Allsworth JE, Stevens E, Macones GA. Short interpregnancy interval: risk of uterine rupture and complications of vaginal birth after caesarean delivery. *Obstet Gynecol*. 2007; 119(5):1075-82.

Standberg-Larsen K, Grønboek M, Andersen AM, Andersen PK, Olsen J. Alcohol drinking pattern during pregnancy and risk of infant mortality. *Epidemiology*. 2009;20(6):884-91.

Standberg-Larsen K, Nielsen NR, Grøenbaek M, Andersen PK, Olsen J, Andersen AM. Binge drinking in pregnancy and risk of fetal death. *Obstet Gynecol*. 2008;111(3):602-9.

Stangl V, Schad J, Gossing G, Borges A, Baumann G, Stangl K. Maternal heart disease and pregnancy outcome: A single-centre experience. *European J Heart Failure*. 2008; 10:855-860.

Stewart AL, Dean ML, Gregorich SE, Brawarsky P, Hass JS. Race/ethnicity, socioeconomic status and the health of pregnant women. *J Health Psychol*. 2007;12(2):285-300.

Stitely ML, Gherman RB. Labor with abnormal presentation and position. *Obstet Gynecol Clin North Am*. 2005;32(2):165-179.

Sturt AS, Dokubo EK, Sint TT. Antiretroviral therapy (ART) for treating HIV infection in ART-eligible pregnant women. *Cochrane Database Syst Rev*. 2010 Mar 17;3:CD008440.

Sundby J, Vangen S. Fetal outcome in severe maternal morbidity: too many stillbirths. *Acta Obstet Gynecol Scand*. 2009;88(3):343-9.

Surkan PJ, Stephansson O, Dickman PW, Cnattingius S. Previous preterm and small-for-gestational-age births and the subsequent risk of stillbirth. *N Engl J Med*. 2004;350(8):777-85.

Tang Y, Ma CX, Cui W, Chang V, Ariet M, Morse SB, Resnick MB, Roth J. The risk of birth defects in multiple births: a population-based study. *Mat Child Health J*. 2006;10(10):75-81.

Tang CH, Wu CS, Lee TH, Hung ST, yang CY, Lee CH, Chu PH. Preeclampsia-eclampsia and the risk of stroke among peripartum in Taiwan. *Stroke*. 2009;40(4):1162-8.

Temple RC, Aldridge V, Stanley K, Murphy HR. Glycaemic control throughout pregnancy and risk of pre-eclampsia in women with type 1 diabetes. *BJOG*. 2006;113(11):1329-32.

Terry R, Westcott J, O'Shea L, Kelly F. Postpartum outcomes in supine delivery by physicians vs. nonsupine delivery by midwives. *J Amer Osteopathic Assn*. 106; 199-202.

Tennant PW, Pearce MS, Bythell M, Rankin J. 20-year survival of children born with congenital anomalies: a population-based study. *Lancet*. 2010;375(9715):649-56.

Thacker SB, Stroup D, Chang M. Continuous electronic heart rate monitoring for fetal assessment during labor. In: *The Cochrane Database of Systematic Reviews*, 2003.

Thanajiraprapa T, Phupong V. Pregnancy complications in women with heart disease. *J Matern Fetal Neonatal Med*. 2009; 1-5.
DOI:10.3109/14767050903410698

Thorngren-Jerneck K, Herbst A. Low 5-minute Apgar Score: A population-based register study of 1 million term births. *Obstet Gynecol.* 2001;98:65-70.

Torloni MR., Betrán AP, Daher S, Widmer M, Dolan SM, Menon R, Bergel E, Allen T, Merialdi M. Maternal BMI and preterm birth: a systematic review of the literature with meta-analysis. *J Maternal-Fetal & Neonatal Medicine*; 2009;22(11):957-70.

Torloni MR, Betrán AP, Horta BL, Nakamura MU, Atallah AN, Moron AF, Valente O. Prepregnancy BMI and the risk of gestational diabetes: a systematic review of the literature with meta-analysis. *Obesity Review.* 2009;10(2):194-203.

Tran SH, Caughey AB, Musci TJ. Meconium-stained amniotic fluid is associated with puerperal infections. *Am J Obstet Gynecol.* 2003;189(3):746-50.

Usta IM, Hobeika EM, Musa AA, Gabriel GE, Nassar AH. Placenta previa-accreta: risk factors and complications. *Am J Obstet Gynecol.* 2005; 193(3 Pt 2):1045-9.

Usta IM, Nassar AH. Advanced maternal age. Part 1: obstetric complications. *Am J Perinatol.* 2008;25(8):521-34. Epub 2008 Sep 4.

Vambergue A, Dognint C, Boulognet A, Rejout M, Beausquet S, Fontaine P. Increasing incidence of abnormal glucose tolerance in women with prior abnormal glucose tolerance during pregnancy: DIAGEST 2 study. *Diabetic Medicine.* 2007;25:58-64.

Vergnano S, Sharland M, Kazembe P, Mwansambo C, Heath PT. Neonatal sepsis: an international perspective. *Arch Dis Child Fetal Neonatal Ed.* 2005;90(3):F220-4.

Villar J, Carroli G, Zavaleta N, Donner A, Wojdyla D, Faundes A et al., Maternal and neonatal individual risks and benefits associated with caesarean delivery: multicentre prospective study. *BMJ.* 2007;335:1025.

Villar J, Carroli G, Khan-Neelofur D, Piaggio G, Gülmezoglu M. Patterns of routine antenatal care for low-risk pregnancy. *Cochrane Database of Systematic Reviews.* 2001. Issue 4. Art. No: CD000934. DOI:10.1002/14651858.CD000934.

Villar J, Carroli G, Wojdyla D, Abalos E, Giordano D, Ba'aqueel H, et al., Preeclampsia, gestational hypertension and intrauterine growth restriction, related or independent conditions? *Am J Obstet Gynecol.* 2006; 194(4):921-31.

Viswanathan M, Hartmann K, Palmieri R, Lux L, Swinson T, Lohr KN, Gartlehner G, Thorp J Jr. The use of episiotomy in obstetrical care: A systematic review. *Evidence Report/Technology Assessment No. 112. AHRQ Publication No. 05-E009-2.* Rockville MD. AHRQ. 2005.

Volante E, Moretti S, Pisani F, Bevilacqua G. Early diagnosis of bacterial infection in the neonate. *J Mat Fetal Neonat Med.* 2004;16 Suppl 2:13-6.

Volmanen P, Valanne J, Alahuhta S. Breast-feeding problems after epidural analgesia for labour: a retrospective cohort study of pain, obstetrical procedures and breast-feeding practices. *Int J Obstet Anesth.* 2004;13(1):25-9.

Volmink J, Siegfried N, van der Merwe L, Brocklehurst P. Antiretrovirals for reducing the risk of mother-to-child transmission of HIV infection. *Cochrane Database of Systematic Reviews*. 2007, Issue 1. Art. No:CD003510. DOI:10.1002/14651858.CD003510.pub2.

Watts DH, Lu M, Thompson B, Tuomala RE, Meyer WA, Mendez H, Rich K, et al. Treatment interruption after pregnancy: effects on disease progression and laboratory findings. *Infectious Diseases in Obstetrics and Gynecology*. 2009; Article ID 45617. DOI:10.1155/2009/456717.

Webb RT. Parental mental illness and fatal birth defects in a national birth cohort. *Psychol Med*. 2008 Oct;38(10):1495-503.

Williams KP, Galerneau F. Intrapartum fetal heart rate patterns in the prediction of neonatal acedemia. *Am J Obstet Gynecol*. 2003;188: 820-3.

Williams KP, Farquharson DF, Bebbington M, Dansereau J, Galerneau F, Wilson RD, Shaw D, Kent N. Screening for fetal well-being in a high-risk pregnant population comparing the nonstress test with umbilical artery Doppler velocimetry: a randomized controlled clinical trial. *Am J Obstet Gynecol*. 2003;188(5):1366-71.

Wilson RE, Alio AP, Kirby RS, Salihu HM. Young maternal age and risk of intrapartum stillbirth. *Arch Gynecol Obstet*. 2008;278(3):231-6.

Wisborg K, Kesmodel U, Henriksen TB, Olsen SF, Secher NJ. Exposure to tobacco smoke in utero and the risk of stillbirth and death in the first year of life. *Am J Epidemiol*. 2001;154(4):322-7 .

Wisner KL. Major depression and antidepressant treatment: impact on pregnancy and neonatal outcomes. *Am J Psychiatry*. 2009; 166(5):557-66.

Whitehead NS, Callaghan W, Johnson C, Williams L. Racial, ethnic, and economic disparities in the prevalence of pregnancy complications. *Matern Child Health J*. 2009;13(2):198-205.

Wolfe EL, Davis T, Guydish J, Delucchi KL. Mortality risk associated with perinatal drug and alcohol use in California. *J Perinatol*. 2005;25(2):93-100.

Wong SF, Chow KM, Ho LC. The relative risk of 'fetal distress' in pregnancy associated meconium-stained liquor at different gestation. *J Obstet Gynaecol*. 2002; 22:594-9.

Wong C, Scavone B, Dugan S, Smith J, Prather H, Ganchiff J, McCarthy R. Incidence of postpartum lumbosacral spine and lower extremity nerve injuries. *Obstet Gynecol*. 2003;101:279-288.

World Health Organization Expert Consultation. Appropriate body mass index for Asian populations and its implication for policy and intervention strategies. *The Lancet*. 2004;363:157-163.

World Health Organization. Make every mother and child count. WHO Geneva. 2005

World Health Organization. Making a difference in countries. Strategic approach to improving maternal and newborn survival and health. WHO Geneva. 2006

Wu S, Kocherginsky M, Hibbard JU. Abnormal placentation: twenty year analysis. *Am J Obstet Gynecol.* 2005;192(5):1458-61.

Xiao R, Sorensen TK, Williams MA, Luthy DA. Influence of pre-eclampsia on fetal growth. *J Matern Fetal Neonatal Med.* 2003;13(3):157-62.

Xu H, Mas-Calvet M, Wei S, Luo Z, Fraser F. Abnormal fetal health rate tracing patterns in patients with thick meconium staining of the amniotic fluid: Association with perinatal outcomes. *Am J Obstet Gynecol.* 2009; 200:283.e1-283.e7.

Yang X, Hsu-Hage B, Zhang H, Zhang C, Zhang Y, Zhang C. Women with impaired glucose tolerance during pregnancy have significantly poorer pregnancy outcomes. *Diabetes Care.* 2002;25(9):1619-24.

Yang Q, Wen SW, Oppenheimer L, Chen XK, Black D, Gao J, Walker MC. Association of caesarean delivery for first birth with placenta praevia and placental abruption in second pregnancy. *BJOG.* 2007;114(5):609-13.

Yang Q, Wen SW, Phillips K, Oppenheimer L, Black D, Walker MC. Comparison of maternal risk factors between placental abruption and placenta previa. *Am J Perinatol.* 2009; 26(4):279-86.

Yokoe DS, Christiansen CL, Johnson R, Sands KE, Livingston J, Shtatland ES, Platt R. Epidemiology of and surveillance for postpartum infections. *Emerg Infect Dis.* 2001;7:837-841.

Yokoe DS, Noskin GA, Cunningham SM, Zuccotti G, Plaskett T, Fraser VJ, Olsen MA, Tokars JI, Solomon S, Perl TM, Cosgrove SE, Tilson RS, Greenbaum M, Hooper DC, Sands KE, Tully J, Herwaldt L, Diekema DJ, Wong ES, Climo M, Platt R. Enhanced identification of postoperative infections among inpatients. *Emerg Infect Dis.* 2004;10(11):1924-30.

Youssef R, Ramalingam U, Macleod M, Murphy DJ. Cohort study of maternal and neonatal morbidity in relation to use of episiotomy at instrumental vaginal delivery. *Br J Obstet Gynaecol.* 2005;112(7):941-5.

Yükesoy G, Ozkan S, Bodur H, Tan T, Calişkan E, Vural B, Corakçi A. Maternal and perinatal outcome in pregnancies complicated with hypertensive disorder of pregnancy: a seven year experience of a tertiary care center. *Arch Gynecol Obstet.* 2005; 273(1):43-9.

Zetterstrom K, Lindeberg SN, Haglund B, Hanson U. Chronic hypertension as a risk factor for offspring to be born small for gestational age. *Acta Obstet Gynecol Scand.* 2006;85(9):1046-50.

Ziadeh SM, Sunna E. Obstetric and perinatal outcome of pregnancies with term labour and meconium-stained amniotic fluid. *Arch Gynecol Obstet.* 2000;264:84-7.

Zhang X, Kramer MS. Variations in mortality and morbidity by gestational age among infants born at term. *J Pediatrics.* 2009; 154(3): 358-62, 262.3.1.

Zhang J, Troendle JF, Levine RJ. Risks of hypertensive disorders in the second pregnancy. *Paediatr Perinat Epidemiol.* 2001 Jul;15(3):226-31

Zhu BP, Le T. Effect of interpregnancy interval on infant low birth weight: a retrospective cohort study using the Michigan Maternally Linked Birth Database. *Matern Child Health J.* 2003;7(3):169-78.

Zwart JJ, Richters A, Ory F, deVries JT, Bloemenkamp KW, van Roosmalen J. Eclampsia in the Netherlands. *Obstet Gynecol.* 2008;112(4):820-7.