

OPTIMALITY INDEX – US

June 2009

REFERENCE CITATIONS

(The evidence in support of the criterion of optimality established for each item in the index)

American College of Obstetricians and Gynecologist (ACOG) Committee on Practice Bulletins. Clinical management guidelines for obstetricians-gynecologists. Prenatal diagnosis of fetal chromosomal abnormalities. *Obstet Gynecol.* 2001;97(5 Pt 1):suppl 1-12.

ACOG Committee on Practice Bulletins. ACOG Practice Bulletin. Chronic hypertension in pregnancy. *Obstet Gynecol.* 2001;98(1):suppl 177-85.

ACOG Committee on Practice Bulletins. Technical Bulletin Number 243. *Int J Gynecol Obstet.* 1998;61:79-86.

ACOG. Prevention of early-onset group B streptococcal disease in newborns. ACOG Committee Opinion #279. 2002.

ACOG. Premature rupture of membranes. Clinical Management Guidelines for Obstetricians-Gynecologists. 1998.

Agency for Health Care Research & Quality (AHRQ). Management of chronic hypertension during pregnancy. Evidence Report/Technology Assessment #14. AHRQ Publication # 00-E0. Rockville, MA: AHRQ, 2000.

Alfirevic Z. Early amniocentesis versus transabdominal chorion villus sampling for prenatal diagnosis. In: *The Cochrane Database of Systematic Reviews*, 2003.

Alfirevic Z, Gosden CM, Neilson JP. Chorion villus sampling versus amniocentesis for prenatal diagnosis In: *The Cochrane Database of Systematic Reviews*, 2003.

Alfirevic Z, Neilson JP. Biophysical profile for fetal assessment in high risk pregnancies. In: *The Cochrane Database of Systematic Reviews*, 2003.

Alfirevic Z, Roberts D, Martlew V. How strong is the association between maternal thrombophilia and adverse pregnancy outcome: A systematic review. *European J Obstet Gynecol & Reprod Biol.* 2002;101:6-14.

Algert C, Morris J, Simpson J, Ford J, Roberts C. Labor before a primary cesarean delivery. *Obstetrics & Gynecology.* 2008;112:1061-6.

Allen L. Anemia and iron deficiency: effects on pregnancy outcome. *Am J Clin Nutr.* 2000;71(Suppl):1280S-4S.

Allen MC, Donohue PK, Dusman AE. The limit of viability - neonatal outcome of infants born at 22-35 weeks gestation. *NEJM.* 1993;329:597.

Alonso A, Soto I, Urgelles MF, Corte JR, Rodriguez MJ, Pinto CR. Acquired and inherited thrombophilia in women with unexplained fetal losses. *Am J Obstet Gynecol.* 2002;187(5):1337-42.

Althaus J, Petersen S, Fox H, Holcroft C, Graham E. Can electronic fetal monitoring identify preterm neonates with cerebral white matter injury? *Obstet Gynecol.* 2005;105(3):458-465.

American Academy of Pediatrics (AAP), ACOG. Use and abuse of the Apgar score. ACOG Committee Opinion 333. Washington, DC, 2006.

AAP, American Heart Association (AHA). Neonatal resuscitation textbook 4th Ed. AHA, Dallas TX, 2000.

AAP. Report of Committee on Infectious Diseases, 24th edition. AAP, Elk Grove, IL, 1997.

Ananth CV, Demissie K, Smulian JC, Vintzileos AM. Placenta previa in singleton and twin births in the United States, 1989 through 1998: a comparison of risk factor profiles and associated conditions. *Am J Obstet Gynecol.* 2003;188(1):275-81.

Ananth CV, Savitz DA. Vaginal bleeding and adverse reproductive outcomes: a meta-analysis. *Paediatr Perinat Epidemiol.* 1994;8(1):62-78.

Ananth CV, Smulian JC, Vintzileos AM. The association of placenta praevia with history of cesarean delivery and abortion: a metaanalysis. *Am J Obstet Gynecol.* 1997;177(5):1071-8.

Ananth CV, Wilcox AJ. Placental abruption and perinatal mortality in the United States. *Am J Epidemiol.* 2001;153(4):332-7.

Anderson GC, Moore E, Hepworth J, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. In: *The Cochrane Database of Systematic Reviews*, 2003.

Andreoli SP. Acute renal failure in the newborn. *Semin Perinatol.* 2004;28(2):112-23.

Arlettaz R, Kashiwagi M, Das-Kundu S, Fauchere JC, Lang A, Bucher JU. Methadone maintenance program in pregnancy in a Swiss perinatal center (II): neonatal outcome and social resources. *Acta Obstet Gynecol Scand.* 2005;84(2):145-50.

Author. Report of the National High Blood Pressure Education Program Working Group on High Blood Pressure in Pregnancy. *Am J Obstet Gynecol.* 2000;183(1):S1-S22.

Aveyard P, Cheng KK, Manaseki S, Gardosi J. The risk of preterm delivery in women from different ethnic groups. *Br J Obstet Gynaecol.* 2002;109(8):894-9.

Bada HS, Das A, Bauer CR, Shankaran S, Lester B, LaGasse L, Hammond J, Wright LL, Higgins R. Impact of prenatal cocaine exposure on child behavior problems through school age. *Pediatrics*. 2007;119(2): e348-59.

Bailey RE. Intrapartum Fetal Surveillance. ALSO Guidelines. Available at:
http://www.aafp.org/online/etc/medialib/aafp_org/documents/cme/courses/clin/also/fetalsurveillance.Par.0001.File.tmp/FetalSurveillance20070830final.pdf

Bailie MD. (Ed) Renal function and disease. *Clin Perinatol*. 1992;19:(entire issue).

Barbosa-Cesnik C, Schwartz K, Foxman B. Lactation mastitis. *JAMA*. 2003;289(13):1609-1612.

Baschat AA, Harman CR. Antenatal assessment of the growth restricted fetus. *Curr Opin Obstet Gynecol*. 2001;13(2):161-8.

Basso O, Christensen K, Olsen J. Higher risk of pre-eclampsia after change of partner. An effect of longer interpregnancy intervals? *J.Epidemiol*. 2001;12(6):624-9.

Beck C. Predictors of postpartum depression: an update. *Nurs Res*. 2001;50(5):275-85.

Ben-Haroush A, Yogev Y, Bar J, Glickman H, Kaplan B, Hod M. Indicated labor induction with vaginal prostaglandin E2 increases the risk of cesarean section even in multiparous women with no previous cesarean section. *J Perinat Med*. 2004;32(1):31-6.

Benson RC, Pernoll ML. *Handbook of Obstetrics & Gynecology*, 9th edition. New York: McGraw-Hill, 1994.

Berg CJ, Chang J, Callaghan WM, Whitehead SJ. Pregnancy-related mortality in the United States, 1991-1997. *Obstet Gynecol*. 2003;101(2):289-96.

Bhatla N, Lal S, Behera G, Kriplani A, Mittal S, Agarwal N, Talwar KK. Cardiac disease in pregnancy. *Int J Gynaecol Obstet*. 2003;82(2):153-9.

Bloom SL, Yost NP, McIntire DD, Leveno KJ. Recurrence of preterm birth in singleton and twin pregnancies. *Obstet Gynecol*. 2001;98(3):379-85.

Blume HK, Li CI, Loch CM, Koepsell TD. Intrapartum fever and chorioamnionitis as risks for encephalopathy in term newborns: a case-control study. *Developmental Medicine and Child Neurology*. 2008;50(1):19-24.

Bouyer J, Coste J, Shojaei T, Pouly JL, Fernandez H, Gerbaud L, Job-Spira N. Risk factors for ectopic pregnancy: a comprehensive analysis based on a large case-control, population-based study in France. *Am J Epidemiol*. 2003;157(3):185-94.

Boy A, Salihu HM. Intimate partner violence and birth outcomes: a systematic review. *Int J Fertil Womens Med*. 2004;49(4):159-64.

- Brabin BJ, Hakimi M, Pelletier D. An analysis of anemia and pregnancy-related maternal mortality. *J Nutr.* 2001;131(2S-2):604S-614S.
- Brady MT. Health care-associated infections in the neonatal intensive care unit. *Am J Infect Control.* 2005;33(5):268-75.
- Branum AM, Schoendorf KC. Changing patterns of low birthweight and preterm birth in the United States, 1981-98. *Paediatr Perinat Epidemiol.* 2002;16(1):8-15.
- Briggs GG, Freeman RK, Yaffe SJ. *Drugs in pregnancy & lactation* 6th Ed. Philadelphia: Williams & Wilkins. 2002.
- Brocklehurst P, Volmink J. Antiretrovirals for reducing the risk of mother-to-child transmission of HIV infection. In: *The Cochrane Database of Systematic Reviews*, 2002.
- Buchmayer SM, Sparen P, Cnattingius S. Previous pregnancy loss: risks related to severity of preterm delivery. *Am J Obstet Gynecol.* 2004;191(3):1225-31.
- Bussel JB. Fetal and neonatal cytopenias: what have we learned? *Am J Perinatol.* 2003;20(8):425-31.
- Cammu H, Martens G, Ruysinck G, Amy JJ. Outcome after elective labor induction in nulliparous women: matched cohort study. *Am J Obstet Gynecol.* 2002; 186: 240-4.
- Canterino JC, Ananth CV, Smulian J, Harrigan JT, Vintzileos AM. Maternal age and risk of fetal death in singleton gestations: USA, 1995-2000. *J Matern Fetal Neonatal Med.* 2004;15(3):193-7.
- Carfoot S, Williamson P, Dickson R. A systematic review of randomized controlled trials evaluating the effect of mother/baby skin-to-skin care on successful breastfeeding. *Midwifery.* 2003;19:148-55.
- Carroli G, Belizan J. Episiotomy for vaginal birth. In: *The Cochrane Database of Systematic Reviews*, 2002.
- Carroli G, Rooney C, Villar J. How effective is antenatal care in preventing maternal mortality and serious morbidity? An overview of the evidence. *Paediatr Perinat Epidemiol.* 2001;15 Suppl 1:1-42.
- Casson, IF, Clarke, CA, Howard, CV, McKendrick, O, Pennycook S, Pharoah PO, Plattt MJ, Stanisstreet M, VanVeszen D, Walkinshaw S. Outcomes of pregnancy in insulin dependent women: results of a five year cohort study. *BMJ.* 1997; 315: 275–278.
- Cederholm M, Haglund B, Axelsson O. Maternal complications following amniocentesis and chorionic villus sampling for prenatal karyotyping. *BJOG.* 2003;110(4):392-9.

- Cederhold M, Haglund B, Axelsson O. Infant morbidity following amniocentesis and chorionic villus sampling for prenatal karyotyping. *BJOG*. 2005;112(4):394-402.
- Centers for Disease Control and Prevention (CDC). Prevention of perinatal Group B strep disease: A public health perspective. *MMWR Morb Mortal Wkly Rep*. 1996;45 (RR7).
- Centers for Disease Control and Prevention. National Task Force on Fetal Alcohol Syndrome and Fetal Alcohol Effect - Defining the national agenda for fetal alcohol syndrome and other prenatal alcohol-related effects. *MMWR Morb Mortal Wkly Rep* 2002;51 (RR14):9-12.
- Chao AS, Huang JY, Lien R, Kung FT, Chen PJ, Hsieh PC. Pregnancy in women who undergo long-term hemodialysis. *Am J Obstet Gynecol*. 2002;187(1):152-6.
- Chalmers B, Porter R. Assessing effective care in normal labor: The Bologna score. *Birth*. 2001;28(2):79-83.
- Chaim W, Bashiri A, Bar-David J, Shoham-Vardi I, Mazor M. Prevalence and clinical significance of postpartum endometritis and wound infection. *Infect Dis Obstet Gynecol*. 2000;8:77-82.
- Chalupczak P, Kolasinska-Kloch W, Jach R, Basta A. Pregnancy in patients with heart disease. *Clin Exp Obstet Gynecol*. 2004;31(4):271-3.
- Charpak N, Ruiz-Pelaez JG, Figueroa de CZ, Charpak Y. A randomized, controlled trial of kangaroo mother care. Results of follow-up at 1 year of corrected age. *Pediatrics*. 2001;108(5):1072-79.
- Chhabra A, Dhorey M. Retained placenta continues to be fatal but frequency can be reduced. *J Obstet Gynecol*. 2002;22(6):630-3.
- Chu SY et al. Gestational weight gain by body mass index among US women delivering live infants, 2004-2005: fueling future obesity. *Am J Obstet Gynecol*. 2009; 271:271e1-271e7.
- Christou H, Brodsky D. Lung injury and bronchopulmonary dysplasia in newborn infants. *J Intensive Care Med*. 2005;20(2):76-87.
- Clausen TD, Mathiesen E, Ekbom P, Hellmuth E, Mandrup-Poulsen T, Damm P. Poor pregnancy outcome in women with type 2 diabetes. *Diabetes Care*. 2005;28(2):323-8.
- Cleary-Goldman J, Malone FD, Vidaver J, Ball RH, Nyberg DA, Comstock CH, Saade GR, Eddleman KA, Klugman S, Dugoff L, Timor-Tritsch IE, Craigo SD, Carr SR, Wolfe HM, Bianchi DW, D'Alton M. Impact of maternal age on obstetric outcome. *Obstet Gynecol*. 2005;105(5 Pt 1):983-90.
- Conde-Aguelo A, Belizan J, Lindmark G. Maternal morbidity and mortality associated with multiple gestations. *Obstet Gynecol*. 2000;95:899-904.

Conde-Agudelo A, Diaz-Rossello JL, Belizan JM. Kangaroo mother care to reduce mortality in low birthweight infants. In: The Cochrane Database of Systematic Reviews, 2003.

Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. JAMA. 2006;295(15):1809-23.

Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: A systematic review. AJOG. 2007;196(4):297-308.

Cotter AM, Martin CM, O'Leary JJ, Daly SF. Increased fetal RhD gene in the maternal circulation in early pregnancy is associated with an increased risk of pre-eclampsia. BJOG. 2005;112(5):584-7.

Crowther CA. Anti-D administration in pregnancy for preventing Rhesus alloimmunisation. In: The Cochrane Database of Systematic Reviews, 2003.

Crowther CA, Hiller JE, Moss JR, McPhee AJ, Jeffries WS, Robinson JS. Effect of treatment of gestational diabetes mellitus on pregnancy outcomes. N Engl J Med. 2005;352(24):2477-86.

Crowther C, Middleton P. Anti-D administration after childbirth for preventing Rhesus alloimmunisation. In: The Cochrane Database of Systematic Reviews, 2003.

Daltveit A, Tollanes MC, Pihlstrom H, Irgens L. Cesarean delivery and subsequent pregnancies. Obstetrics & Gynecology. 2008; 111(6):1327-34.

Dasgupta SJ, Gill AB. Hypotension in the very low birthweight infant: the old, the new, and the uncertain. Arch Dis Child Fetal Neonatal Ed. 2003;88(6):F450-4.

Dashe JS, Sheffield JS, Olscher DA, Todd SJ, Jackson GL, Wendel GD. Relationship between maternal methadone dosage and neonatal withdrawal. Obstet Gynecol. 2002;100(6):1244-9.

Decca L, Daldoss C, Fratelli N, Lojaco A, Slompo M, Stegher C, Valcamonico A, Frusca T. Labor course and delivery in epidural analgesia: a case-control study. J Matern Fetal Neonatal Med. 2004;16(2):115-8.

Demissie K, Rhoads GG, Smulian JC, Balasubramanian BA, Gandhi K, Joseph KS, Kramer M. Operative vaginal delivery and neonatal and infant adverse outcomes: population based retrospective analysis. BMJ. 2004;329(7456):24-9.

Donnelly A, Snowden HM, Renfrew MJ, Woolridge MW. Commercial hospital discharge packs for breastfeeding women. In: The Cochrane Database of Systematic Reviews, 2005.

Dyson L, McCormick F, Renfrew MJ. Interventions for promoting the initiation of breastfeeding. In: The Cochrane Database of Systematic Reviews, 2005.

Ekeroma AJ, Ansari A, Stirrat GM. Blood transfusion in obstetrics and gynecology. *Br J Obstet Gynaecol.* 1997;104:278-284.

Elbourne D, Wiseman RA. Types of intra-muscular opioids for maternal pain relief in labour. In: *The Cochrane Database of Systematic Reviews*, 2003

Eyler FD, Behnke M, Conlon M, Woods N, Wobie K.. Birth outcome from a prospective, matched study of prenatal crack/cocaine use: I. Interactive and dose effects on health and growth. *Pediatrics.* 1998;101(2):229-37.

Ezra Y, Michaelson-Cohen R, Abramov Y, Rojansky N. Prelabor rupture of the membranes at term: when to induce labor? *Eur J Obstet Gynecol Reprod Biol.* 2004;115(1):23-7.

Ferrer RL, Sibai BM, Mulrow CD, Chiquette E, Stevens KR, Cornell J. Management of mild chronic hypertension during pregnancy: a review. *Obstet Gynecol.* 2000;96(5 Pt 2):849-60.

Fischer MJ, Lehnerz SD, Hebert JR, Parikh CR. Kidney disease is an independent risk factor for adverse fetal and maternal outcomes in pregnancy. *Am J Kidney Dis.* 2004;43(4):415-23.

Fleming VE, Hagen S, Niven C. Does perineal suturing make a difference? The SUNS trial. *Br J Obstet Gynaecol.* 2003;119(7):684-9

Flenady V, King J. Antibiotics for prelabor rupture of membranes at or near term. In: *The Cochrane Database of Systematic Reviews*, 2003

Forna F, Jamieson DJ, Sanders D, Lindsay MK. Pregnancy outcomes in foreign-born and US-born women. *Int J Gynaecol Obstet.* 2003;83(3):257-65.

Foxman B. Epidemiology of urinary tract infections: incidence, morbidity and economic costs. *Am J Med.* 2002;113 Suppl 1A:5S-13S.

French L. Prevention and treatment of postpartum endometritis. *Curr Women's Health Rep.* 2003;3(4):274-9.

French LM, Smaill FM. Antibiotic regimens for endometritis after delivery. In: *The Cochrane Database of Systematic Reviews*, 2002.

Frias AE Jr, Luikenaar RA, Sullivan AE, Lee RM, Porter TF, Branch DW, Silver RM. Poor obstetric outcome in subsequent pregnancies in women with prior fetal death. *Obstet Gynecol.* 2004;104(3):521-6.

Froen JF, Gardosi JO, Thurmann A, Francis A, Stray-Pedersen B. Restricted fetal growth in sudden intrauterine unexplained death. *Acta Obstet Gynecol Scand.* 2004;83(9):801-7.

Gabbe SG, Niebyl JR, Simpson JL. *Obstetrics. Normal and problem pregnancies.* 4th Ed. New York: Churchill Livingstone, 2002.

Getahun D, Amre D, Rhoads GG, Demissie K. Maternal and obstetric risk factors for sudden infant death syndrome in the United States. *Obstet Gynecol.* 2004;103(4):646-52.

Gourounti K, Sandall J. Admission cardiotocography versus intermittent auscultation of fetal heart rate: effects on neonatal Apgar score, on the rate of caesarean sections and on the rate of instrumental deliver – a systematic review. *Int J Nursing Stud.* 2007;44(6):1029-35.

Granelli SL, McGrath JM. Neonatal seizures: diagnosis, pharmacologic interventions, and outcomes. *J Perinat Neonatal Nurs.* 2004;18(3):275-87.

Gregory KD, Korst LM. Age and racial/ethnic differences in maternal, fetal, and placental conditions in laboring patients. *Am J Obstet Gynecol.* 2003;188(6):1602-6.

Grobman W, Lai Y, Landon MB, et al. Can a prediction model for vaginal birth after cesarean also predict the probability of morbidity related to a trial of labor ? *Am J Obstet Gynecol.* 2009;200:56.e1-56.e6

Gul A, Cebeci A, Aslan H, Polat I, Ozdemir A, Ceylan Y. Perinatal outcomes in severe preeclampsia-eclampsia with and without HELLP syndrome. *Gynecol Obstet Invest.* 2005;59(2):113-8.

Gupta JK, Nikodem VC. Woman's position during second stage of labour. In: *The Cochrane Database of Systematic Reviews*, 2000.

Gupta JK, Hofmeyr GR. Position for women during second stage of labour. In: *The Cochrane Database of Systematic Reviews*, 2003.

Habek D. Fetal shoulder dystocia. *Acta Med Croatica.* 2002;56:57-63.

Hansen A, Wisborg K, Uldbjerg N, Henriksen TB. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *British Medical Journal.* 2008; 336:85-87.

Hanson S, Clark S, Foster J. Active pushing vs passive fetal descent in the second state of labor: A randomized controlled trial. *Obstet Gynecol.* 2002;99:29-34.

Hartmann K, Viswanathan M, Palmieri R, Gertlehner G, Thorp J, Lohr KN. Outcomes of routine episiotomy: a systematic review. *JAMA.* 2005;293:2141-8.

Heinonen S, Kirkinen P. Pregnancy outcome after previous stillbirth resulting from causes other than maternal conditions and fetal abnormalities. *Birth.* 2000;27(1):3307.

Henriksen TB, Hjollund NH, Jensen TK, Bonde JP, Andersson AM, Kolstad H, Ernst E, Giwercman A, Skakkebaek NE, Olsen J. Alcohol consumption at the time of conception and spontaneous abortion. *Am J Epidemiol.* 2004;160(7):661-7.

Herbst MA, Mercer BM, Beazley D, Meyer N, Carr T. Relationship of prenatal care and perinatal morbidity in low-birth-weight infants. *Am J Obstet Gynecol.* 2003;189(4):930-3.

Hill JB, Alexander JM, Sharma SK, McIntire DD, Leveno KJ. A comparison of the effects of epidural and meperidine analgesia during labor on fetal heart rate. *Obstet Gynecol.* 2003;102(2):333-7.

Hill JB, Sheffield JS, McIntire DD, Wendel GD Jr. Acute pyelonephritis in pregnancy. *Obstet Gynecol.* 2005;105(1):18-23.

HIV/AIDS Treatment Information Service Available on-line at: <http://www.hivatis.org>

Hnat MD, Sibai BM, Caritis S, Hauth J, Lindheimer MD, MacPherson C, VanDorsten JP, Landon M, Miodovnik M, Paul R, Meis P, Thurnau G, Dombrowski M; National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. Perinatal outcome in women with recurrent preeclampsia compared with women who develop preeclampsia as nulliparas. *Am J Obstet Gynecol.* 2002;186(3):422-6.

Hoch M, Netz H. Heart failure in pediatric patients. *Thorac Cardiovasc Surg.* 2005;53 Suppl 2:S129-34.

Hodnett ED. Continuity of caregivers for care during pregnancy and childbirth In: *The Cochrane Database of Systematic Reviews*, 2003.

Hodness ED, Gates S, Hofmeyr GJ, Sakala A. Continuous support for women during childbirth. *The Cochrane Database of Systematic Reviews.* 2007; Issue 3.

Hoffman JA, Mason EO, Schutze GE, Tan TQ, Barson WJ, Givner LB, Wald ER, Bradley JS, Yogev R, Kaplan SL. Streptococcus pneumoniae infections in the neonate. *Pediatrics.* 2003;112(5):1095-102.

Holzman C, Lin X, Senagore P, Chung H. Histologic chorioamnionitis and preterm delivery. *Am J Epidemiol.* 2007;166(7):786-94.

Hordnes K, Bergsjø P. Severe lacerations after childbirth. *Acta Obstet Gynecol.* 1993;72:413-422.

Howard LM, Goss C, Leese M, Appleby L, Thornicroft G. The psychosocial outcome of pregnancy in women with psychotic disorders. *Schizophr Res.* 2004;71(1):49-60.

Howard LM, Goss C, Leese M, Thornicroft G. Medical outcome of pregnancy in women with psychotic disorders and their infants in the first year after birth. *Br J Psychiatry.* 2003;182:63-7.

Howell CJ. Epidural versus non-epidural analgesia for pain relief in labour. In: *The Cochrane Database of Systematic Reviews*, 2003.

- Howell CJ, Dean T, Lucking L, Dziedzic K, Jones, PW, Johanson RB. Randomised study of long term outcome after epidural versus non-epidural analgesia during labor. *BMJ*. 2002; 325(7360):357.
- Hurd YL, Wang X, Anderson V, Beck O, Minkoff H, Dow-Edwards D. Marijuana impairs growth in mid-gestation fetuses. *Neurotoxicol Teratol*. 2005;27(2):221-9.
- Ibison JM. Ethnicity and mode of delivery in “low-risk” first time mothers. *Eur J Obstet Gynecol Reprod Biol*. 2005;118(2):199-205.
- Ingemarsson I. Controversies: Premature rupture of membranes at term - no advantage of delaying induction > 24 hours. *J Perinat Med*. 1996;24:573-579.
- Jablensky AV, Morgan V, Zubrick SR, Bower C, Yellachich LA. Pregnancy, delivery, and neonatal complications in a population cohort of women with schizophrenia and major affective disorders. *Am J Psychiatry*. 2005;162(1):79-91.
- Jacobsson B, Ladfors L, Milsom I. Advanced maternal age and adverse perinatal outcome. *Obstet Gynecol*. 2004;104(4):727-33.
- Jain S, Arya VK, Gopalan S, Jain V. Analgesic efficacy of intramuscular opioids versus epidural analgesia in labor. *Int J Gynaecol Obstet*. 2003;83(1):19-27.
- Johanson RB, Menon V. Vacuum extraction versus forceps for assisted vaginal delivery. In: *The Cochrane Database of Systematic Reviews*, 2005.
- Jonsson S, Pulkkinen MO. Mastitis today: Incidence, prevention and treatment. *Ann Chir Gynaecol*. 1994;208(suppl):84-87.
- Kahana B, Sheiner E, Levy A, Lazer S, Mazor M. Umbilical cord prolapse and perinatal outcome. *Int J Gynaecol Obstet*. 2004;84(2):127-32.
- Kalil A, Kunz J. Teenage childbearing, marital status, and depressive symptoms in later life. *Child Dev* 2002;73(6):1748-60.
- Kassanos D, Siristatidis C, Vitoratos N, Salamalekis E, Creatsas G. The clinical significance of Doppler findings in fetal middle cerebral artery during labor. *Eur J Obstet Gynecol Reprod Biol*. 2003;109(1):45-50.
- Kaye DK, Mirembe FM, Bantebya G, Johannsson A, Eksrom AM. Domestic violence during pregnancy and risk of low birthweight and maternal complications: a prospective cohort study at Mulago Hospital, Uganda. *Trop med Int Health*. 2006;11(10):1576-84.
- Keirse MJ, Ottervanger H, Smit W. Controversies: Prelabor rupture of the membranes at term: the case for expectant management. *J Perinat Med*. 1996;24:563-572.
- Kesmodel U, Wisborg K, Olsen SF, Henriksen TB, Secher NJ. Moderate alcohol intake in pregnancy and the risk of spontaneous abortion. *Alcohol*. 2002;37(1):87-92.

- Kesmodel U, Wisborg K, Olsen SF, Henriksen TB, Secher NJ. Moderate alcohol intake during pregnancy and the risk of stillbirth and death in the first year of life. *Am J Epidemiol.* 2002;155(4):305-12.
- Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PFA. WHO analysis of causes of maternal death: a systematic review. *Lancet.* 2006;367:1066-1074.
- Kinsella JP, Abman SH. Recent developments in the pathophysiology and treatment of persistent pulmonary hypertension of the newborn. *J Pediatrics.* 1995;126:853.
- Kliegman RM, Walker WA, Yolken RH. Necrotizing enterocolitis: Research agenda for a disease of unknown etiology and pathogenesis. *Pediatr Res.* 1993;34:701-708.
- Koike T, Minakami H, Izumi A, Watanabe T, Matsubara S, Sato I. Recurrence risk of preterm birth due to preeclampsia. *Gynecol Obstet Invest.* 2002;53(1):22-7.
- Konduri GG. New approaches for persistent pulmonary hypertension of newborn. *Clin Perinatol.* 2004;31(3):591-611.
- Kristensen J, Vestergaard M, Wisborg K, Kesmodel U, Secher NJ. Pre-pregnancy weight and the risk of stillbirth and neonatal death. *BJOG.* 2005;112(4):403-8.
- Kuban KCK. Intracranial Hemorrhage. In Cloherty JP, Stark AR (Eds.). *Manual of Neonatal Care*, 4th Edition. Philadelphia: Lippincott-Raven, 1998.
- Langer O, Yogeve Y, Most O, Xenakis EM. Gestational diabetes: the consequences of not treating. *Am J Obstet Gynecol.* 2005;192(4):989-97.
- Larma AJ, Silva A, Holcroft C, Thompson R, Donohue P, Graham E. Intrapartum electronic fetal heart rate monitoring and the identification of metabolic acidosis and hypoxic-ischemic encephalopathy. *Am J Obstet Gynecol.* 2007;197:301e1-e8.
- Lasker JN, Coyle B, Li K, Ortynsky M. Assessment of risk factors for low birth weight deliveries. *Health Care Women Int.* 2005;26(3):262-80.
- Lemons JA, Bauer CR, Oh W, Korones SP, Papile LA, Stoll BJ, Verter J, Temprosa M, Wright HL, Ehrenkranz RA, Fanaroff AA, Stark A, Carlo W, Tyson JE, Donovan EF, Shankaran S, Stevenson D. Very low birth weight outcomes of the National Institute of Child Health and Human Development Neonatal Research Network, January 1995 through December 1996. *Pediatrics.* 2001;107(1):E1.
- Lepercq J, Coste J, Theau A, Dubois-Laforgue D, Timsit J. Factors associated with preterm delivery in women with type 1 diabetes: a cohort study. *Diabetes Care.* 2004;27(12):2824-8.

- Lieber CS. Molecular basis and metabolic consequences of ethanol metabolism. In: Heather N, Peters, TJ and Stockwell T, Eds, International handbook of alcohol dependence and problems, Wiley & Sons, West Sussex, England, 2001.
- Lin C, Raynor BD. Risk of uterine rupture in labor induction of patients with prior cesarean section: an inner city hospital experience. *Am J Obstet Gynecol.* 2004;190(5):1476-8.
- Liu BY, Wang CC, Lau TK, Chu CY, Phil M, Pang CP, Rogers MS, Leung TN. Meconium-stained liquor during labor is associated with raised neonatal cord blood 8-iso-prostaglandin F₂α concentration. *Am J Obstet Gynecol.* 2005;192(1):289-94.
- Locatelli A, Regalia AL, Patregnani C, Ratti M, Toso L, Ghidini A. Prognostic value of change in amniotic fluid color during labor. *Fetal Diagn Ther.* 2005;20(1):5-9.
- Locksmith G, Duff P. Infection, antibiotics and preterm birth. *Semin Perinatol.* 2001;25(5):295-309.
- Lone FW, Qureshi RN, Emanuel F. Maternal anaemia and its impact on perinatal outcome. *Trop Med Int Health.* 2004; 9(4):486-90.
- Ludlow JP, Evans SF, Hulse G. Obstetric and perinatal outcomes in pregnancies associated with illicit substance abuse. *Aust N Z J Obstet Gynaecol.* 2004;44(4):302-6.
- Lundquist M, Olsson A, Nissen E, Norman M. Is it necessary to suture all lacerations after a vaginal delivery? *Birth.* 2000;27(2):79-85.
- Luo ZC, Wilkins R, Kramer MS. Disparities in pregnancy outcomes according to marital and cohabitation status. *Obstet Gynecol.* 2004;103(6):1300-7.
- Macones GA, Cahill A, Pare E, Stamilio DM, Ratcliffe S, Stevens E, Sammel M, Peipert J. Obstetric outcomes in women with two prior cesarean deliveries: is vaginal birth after cesarean delivery a viable option? *Am J Obstet Gynecol.* 2005;192(4):1223-8.
- Magann EP, Channan SP, Bofill JA, Waddell D, Rust OA, Morrison JC. Maternal morbidity and mortality associated with intrauterine fetal demise: five-year experience in a tertiary referral hospital. *South Med J.* 2001;94(5):493-5.
- Many A, Elad R, Yaron Y, Eldor A, Lessing JB, Kupferminc MJ. Third-trimester unexplained intrauterine fetal death is associated with inherited thrombophilia. *Obstet Gynecol.* 2002;99(5 Pt 1):684-7.
- Massad LS, Springer G, Jacobson L, Watts H, Anastos K, Korn A, Cejtin H, Stek A, Young M, Schmidt J, Minkoff H. Pregnancy rates and predictors of conception, miscarriage and abortion in US women with HIV. *AIDS.* 2004;18(2):281-6.
- McCowan LM, Buist RG, North RA, Gamble G. Perinatal morbidity in chronic hypertension. *Br J Obstet Gynaecol.* 1996;103(2):123-9.

McDonald S, Prendiville WJ, Elbourne D. Prophylactic syntometrine versus oxytocin for delivery of the placenta. In: The Cochrane Database of Systematic Reviews, 2002.

Meyer KA, Williams P, Hernandez-Diaz S, Cnattingius S. Smoking and the risk of oral clefts: exploring the impact of study designs. *Epidemiology*. 2004;15(6):671-8.

Millar LK, DeBuque L, Wing DA. Uterine contraction frequency during treatment of pyelonephritis in pregnancy and subsequent risk of preterm birth. *J Perinat Med*. 2003;31(1):41-6.

Moore ER, Anderson GC, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. The Cochrane Database of Systematic Reviews. 2007; Issue 4.

Moore LL, Bradless ML, Singer MR, Rothman KJ, Milunsky A. Chromosomal anomalies among the offspring of women with gestational diabetes. *Am J Epidemiol*. 2002;155(8):719-24.

Moroni G, Quaglini S, Banfi G, Caloni M, Finazzi S, Ambroso G, Como G, Ponticelli C. Pregnancy in lupus nephritis. *Am J Kidney Dis*. 2002;40(4):713-20.

Mortensen JT, Thulstrup AM, Larsen H, Moller M, Sorensen HT. Smoking, sex of the offspring, and risk of placental abruption, placenta previa, and preeclampsia: a population-based cohort study. *Acta Obstet Gynecol Scand*. 2001;80(10):894-8.

Murphy DJ, MacKenzie IZ. The mortality and morbidity associated with umbilical cord prolapse. *Br J Obstet Gynaecol*. 1995;102:826-830.

Myers-Helfgott MG, Helfgott AW. Routine use of episiotomy in modern obstetrics: should it be performed? *Obstet Gynecol Clin North Amer*. 1999;26(2):302-325.

Nabet C, Ancel PY, Burguet A, Kaminski M. Smoking during pregnancy and preterm birth according to obstetric history: French national perinatal surveys. *Paediatr Perinat Epidemiol*. 2005;19(2):88-96.

Natale R, Dodman N. Birth can be a hazardous journey: electronic fetal monitoring does not help. *J Obstet Gynaecol Can*. 2003;25(12):1007-9.

National Center for Health Statistics. 1999. cited at <http://www.marchofdimes.com/about us/1529.asp> (accessed 4/8/2003).

National Institute of Child Health and Human Development. Electronic fetal heart rate monitoring: research guidelines for interpretation. National Institute of Child Health and Human Development Research Planning Workshop. *Am J Obstet Gynecol*. 1997;177(6):1385-90.

Ness RB, Grisso JA, Hirschinger N, Markovic N, Shaw LM, Day NL, Kline J. Cocaine and tobacco use and the risk of spontaneous abortion. *N Engl J Med*. 2002;340(5):333-339.

Normand MC, Damato EG. Postcesarean infection. *J Obstet Gynecol Neonat Nurs*. 2001;30:642-648.

Nystedt A, Edvardsson D, Willman A. Epidural analgesia for pain relief in labour and childbirth – a review with a systematic approach. *J Clin Nurs*. 2004;13(4):455-66.

O’Connell MP, Hussain J, MacLennan FA, Lindow SW. Factors associated with a prolonged second state of labour – a case-controlled study of 364 nulliparous labours. *J Obstet Gynaecol*. 2003;23(3):255-7.

Ogunyemi D, Hernandez-Loera GE. The impact of antenatal cocaine use on maternal characteristics and neonatal outcomes. *J Matern Fetal Neonatal Med*. 2004;15(4):253-9.

Olsen O, Jewell MD. Home versus hospital birth. In: *The Cochrane Database of Systematic Reviews*, 2003.

Oron T, Sheiner E, Shoham-Vardi I, Mazor M, Katz M, Hallak M. Risk factors for antepartum fetal death. *J Repro Med*. 2001;46(9):825-30.

Osrin D, Vergnano S, Costello A. Serious bacterial infections in newborn infants in developing countries. *Curr Opin Infect Dis*. 2004;17(3):217-24.

Owens R. Intraventricular hemorrhage in the premature neonate. *Neonatal Netw*. 2005;24(3):55-71.

PAHO. Quantifying the benefits of breastfeeding – A summary of the evidence., 2002. Available on-line at: http://www.paho.org/English/HPP/HPN/benefits_of_BF.htm.

Pang JWY, Heffelfinger JD, Huang GJ, Benedetti TJ, Weiss NS. Outcomes of planned home births in Washington state 1989-1996. *Obstet Gynecol*. 2002;100:253-9.

Parazzini F, Chatenoud L, Surace M, Tozzi L, Salerio B, Bettoni G, Benzi G. Moderate alcohol drinking and risk of preterm birth. *Eur J Clin Nutr*. 2003;57(10):1345-9.

Parnell C, Langhoff-Roos J, Iverson R, Damgaard P. Pushing method in the expulsive phase of labor: A randomized controlled trial. *Acta Obstet Gynecol Scand*. 1993;72:31-35.

Phipps MG, Blume JD, DeMonner SM. Young maternal age associated with increased risk of postneonatal death. *Obstet Gynecol*. 2002; 100(3):481-6.

Propst AM, Thorp JM. Traumatic vulvar hematomas: Conservative versus surgical management. *Southern Med J*. 1998;9:144-146.

- Rasch V. Cigarette, alcohol, and caffeine consumption: risk factors for spontaneous abortion. *Acta Obstet Gynecol Scand.* 2003;82(2):182-8.
- Rasmussen S, Albrechtsen S, Dalaker K. Obstetric history and the risk of placenta previa. *Acta Obstet Gynecol Scand.* 2000;79(6):502-7.
- Rasmussen K. Is there a causal relationship between iron deficiency or iron-deficiency anemia and weight at birth, length of gestation and perinatal mortality? *J Nutr.* 2001;131(2S-2):590S-601S.
- Rasmussen S, Irgens LM, Dalaker K. Outcomes of pregnancies subsequent to placental abruption: a risk assessment. *Acta Obstet Gynecol Scand.* 2000;79(6):496-501.
- Ray JG, Burrows RF, Burrows EA, Vermeulen MJ. MOS HIP: McMaster outcome study of hypertension in pregnancy. *Early Hum Dev.* 2001; 64(2):129-43.
- Reynolds F, Sharma SK, Seed PT. Analgesia in labour and fetal acid-base balance: a meta-analysis comparing epidural with systemic opioid analgesia. *Br J Obstet Gynaecol.* 2002;109:1344-1353.
- Richardson GA, Goldschmidt L, Larkby C. Effects of prenatal cocaine exposure on growth: a longitudinal analysis. *Pediatrics.* 2007; 120(4): e1017-1027.
- Richter R, Bergmann R, Dudenhausen J. Previous caesarean or vaginal delivery: Which mode is a greater risk of perinatal death at the second delivery? *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 2007;132:51-57.
- Ridgeway JJ, Weyrich DL, Benedetti TJ. Fetal heart rate changes associated with uterine rupture. *Obstet Gynecol.* 2004;103(3):506-12.
- Roberts JE. The “push” for evidence: management of the second stage. *J Midwif Womens Health.* 2002;47(1):2-15.
- Rogers J, Wood J, McCandish R, Ayers S, Truesdale A, Elbourne D. Active versus expectant management of third stage of labour: The Hinchingsbrooke randomized controlled trial. *The Lancet.* 1998;351:693-699.
- Rojas MA, Kaplan M, Quevedo M, Sherwonit E, Foster LB, Ehrenkranz RA, Mayes L. Somatic growth of preterm infants during skin-to-skin care versus traditional holding: a randomized, controlled trial. *J Dev Behav Pediatr.* 2003;24(3):163-8.
- Ronsmans C, Graham W, The Lancet Maternal Survival Series Steering Group. Maternal mortality: who, when, where, and why. *Lancet.* 2006;368:1189-200.
- Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskiene E, Samuelsson U. Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood. *Diabetes Metab Res Rev.* 2004;20(2):150-7.
- Samselle C, Hines S. Spontaneous pushing during birth. *J Nurse Midwif.* 1999;33(1):36-39.

Scanlon KS, Yip R, Schieve LA, Cogswell ME. High and low hemoglobin levels during pregnancy: differential risks for preterm birth and small for gestational age. *Obstet Gynecol.* 2000;95 (5 Pt 1):741-8.

Schrag S, Schuchat A. Prevention of neonatal sepsis. *Clin Perinatol.* 2005;32(3):601-15.

Seward PG, Hannah ME, Myhr TL, Farine D, Ohlsson A, Wang EE, Hodnett E, Haque K, Weston JA, Phel G. International multicenter term PROM study: evaluation of predictors of neonatal infection in infants born to patients with premature rupture of membranes at term. *Am J Obstet Gynecol.* 1998; 635-9.

Sharps P, Laughon K, Giangrande S. Intimate partner violence and the childbearing year: Maternal and infant health consequences. *Trauma, Violence and Abuse.* 2007;6(2):105-116.

Shaw AK, Infante-Rivard C, Morrison HI. Use of medication during pregnancy and risk of childhood leukemia (Canada). *Cancer Causes Control.* 2004;15(9):931-7.

Sheiner E, Shoham-Vardi I, Hallak M, Hershkowitz R, Katz M, Mazor M. Placenta previa: obstetric risk factors and pregnancy outcome. *J Mat Fetal Med.* 2001;10(6):414-9.

Sheiner E, Shoham-Vardi I, Hadar A, Hallak M, Hackmon R, Mazor M. Incidence, obstetric risk factors and pregnancy outcome of preterm placental abruption: a retrospective analysis. *J Mat Fetal Neonat Med.* 2002;11(1):34-9.

Sheiner E, Shohan-Vardi I, Hadar A, Hershkowitz R, Sheiner EK, Mazor M. Accidental out-of-hospital delivery as an independent risk factor for perinatal mortality. *J Reprod Med.* 2002; 47(8):625-30.

Shetty A, Stewart K, Stewart G, Rice P, Danielian P, Templeton A. Active management of term prelabor rupture of membranes with oral misoprostol. *Br J Obstet Gynaecol.* 2002, 109: 1354-8.

Shipp T, Zelop C, cohen A, Repke J, Lieberman E. Post-cesarean delivery fever and uterine rupture in a subsequent trial of labor. *Obstet Gynecol.* 2003;101(1):136-139.

Sibai BM, Lindheimer M, Hauth J, Caritis S, VanDorsten P, Klebanoff M, MacPherson C, Landon M, Miodovnik M, Paul R, Meis P, Dombrowshi M. Risk factors for preeclampsia, abruptio placentae, and adverse neonatal outcomes among women with chronic hypertension. *N Engl J Med.* 1998;339:667-671.

Sikorski J, Renfrew MJ, Pindorial S, Wade A. Support for breastfeeding mothers. In: *The Cochrane Database of Systematic Reviews*, 2005.

Silva Idos S, Higgins C, Swerdlow AJ, Laing SP, Slater SD, Pearson DW, Morris AD. Birthweight and other pregnancy outcomes in a cohort of women with pre-gestational insulin-treated diabetes mellitus, Scotland, 1979-95. *Diabet Med.* 2005;22(4):440-7.

Silverman JG, Decker MR, Reed E, Raj A. Intimate partner violence victimization prior to and during pregnancy among women residing in 26 states: associations with maternal and neonatal health. *Am J Obstet Gynecol.* 2006;195(1):140-8.

Simpson LL. Maternal medical disease: risk of antepartum fetal death. *Semin Perinatol.* 2002;26(1):42-50.

Simpson KR, James DC. Effects of immediate versus delayed pushing during second-stage labor on fetal well-being: a randomized clinical trial. *Nurs Res.* 2005;54(3):149-57.

Siu S, Colman J, Sorensen S, Smallhorn J, Farine D, Amankwah K, Spears J, Sermer M. Adverse neonatal and cardiac outcomes are more common in pregnant women with cardiac disease. *Circulation.* 2002;105:2179-2184.

Siu S, Sermer M, Colman J, Alvarez AN, Mercier L, Morton B, Kells C, Bergin M, Kiess M, Marcott F, Taylor D, Gordon E, Spears J, Tam J, Amankwah K, Smallhorn J, Farine D, Sorensen S. Prospective multicenter study of pregnancy outcomes in women with heart disease. *Circulation.* 2001;104:515-521.

Sloan N, Jordan E, Winikoff M. Effects of iron supplementation on maternal hematologic status in pregnancy. *Am J Public Health.* 2002;92(2):268-293.

Smith GC, Pell JP. Teenage pregnancy and risk of adverse perinatal outcomes associated with first and second births: population based retrospective cohort study. *BMJ.* 2001, 323(7326): 1428-9.

Smith GC, Pell JP, Dobbie R. Interpregnancy interval and risk of preterm birth and neonatal death: a retrospective cohort study. *BMJ.* 2003;327(7410):313. *Erratum in* *BMJ* 2003;327(7419):851.

Smulian JC, Potash SK, Lai YL, Scorza WE. Appropriateness of antibiotic use in the postpartum period. *J Mat Fetal Med.* 2001;10:312-317.

Smulian JC, Ananth CV, Vintzileos AM, Scorza WE, Knuppel RA. Fetal deaths in the United States. Influence of high-risk conditions and implications for management. *Obstet Gynecol.* 2002;100(6):1183-9.

Smyth RMD, Alldred SK, Markham C. Amniotomy for shortening spontaneous labour. *Cochrane Database of Systematic Reviews.* 2007; Issue 4.

Soll RF. Synthetic surfactant for respiratory distress syndrome in preterm infants. In: *The Cochrane Database of Systematic Reviews*, 2005.

Soong B, Barnes M. Maternal position in midwife-attended birth and perinatal trauma: Is there an association? *Birth.* 2005;32(3):164-169.

Stephansson O, Dickman PW, Johansson A, Cnattingius S. Maternal weight, pregnancy weight gain and the risk of antepartum stillbirth. *Am J Obstet Gynecol.* 2001;184(3):462-9.

Stitely ML, Gherman RB. Labor with abnormal presentation and position. *Obstet Gynecol Clin North Am.* 2005;32(2):165-179.

Surkan PJ, Stephansson O, Dickman PW, Cnattingius S. Previous preterm and small-for-gestational-age births and the subsequent risk of stillbirth. *N Engl J Med.* 2004;350(8):777-85.

Tang Y, Ma CX, Cui W, Chang V, Ariet M, Morse SB, Resnick MB, Roth J. The risk of birth defects in multiple births: a population-based study. *Mat Child Health J.* 2006;10(10):75-81.

Temple R, Aldridge V, Greenwood R, Heyburn P, Sampson M, Stanley K. Association between outcome of pregnancy and glycaemic control in early pregnancy in type 1 diabetes: population based study. *BMJ.* 2002;325(7375):1275-6.

Thacker SB, Stroup D, Chang M. Continuous electronic heart rate monitoring for fetal assessment during labor. In: *The Cochrane Database of Systematic Reviews*, 2003.

Thompson A. Pushing techniques in the second stage of labour. *J Adv Nurs.* 1993;14:101-105.

Thorngren-Jerneck K, Herbst A. Low 5-minute Apgar Score: A population-based register study of 1 million term births. *Obstet Gynecol.* 2001;98:65-70.

Torloni MR et al. Prepregnancy BMI and the risk of gestational diabetes: a systematic review of the literature with meta-analysis. *Obesity Reviews.* 2009; 10:194-203

Tran SH, Caughey AB, Musci TJ. Meconium-stained amniotic fluid is associated with puerperal infections. *Am J Obstet Gynecol.* 2003;189(3):746-50.

U.S. Department of Health and Human Services. *Smoking and Women's Health: A Report of the Surgeon General.* Atlanta, GA: USDHHS, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2001.

Vahratian A, Zhang J, Troendle JF, Sciscione AC, Hoffman MK. Labor progression and risk of cesarean delivery in electively induced nulliparas. *Obstet Gynecol.* 2005;105(4):698-704.

Varner MW, Esplin MS. Current understanding of genetic factors in preterm birth. *BJOG.* 2005;112 Suppl 1:28-31.

Vergnano S, Sharland M, Kazembe P, Mwansambo C, Heath PT. Neonatal sepsis: an international perspective. *Arch Dis Child Fetal Neonatal Ed.* 2005;90(3):F220-4.

- Villar J, Carroli G, Khan-Neelofur D, Piaggio G, Gülmezoglu M. Patterns of routine antenatal care for low-risk pregnancy. *Cochrane Database of Systematic Reviews*. 2001. Issue 4. Art. No: CD000934. DOI:10.1002/14651858.CD000934.
- Vintzileos AM, Nochimson DJ, Guzman ER, Knuppel RA, Lake M, Schiffrin BS. Intrapartum electronic fetal heart rate monitoring versus intermittent auscultation: a meta-analysis. *Obstet Gynecol*. 1995;85(1):149-55.
- Viswanathan M, Hartmann K, Palmieri R, Lux L, Swinson T, Lohr KN, Gartlehner G, Thorp J Jr. The use of episiotomy in obstetrical care: A systematic review. Evidence Report/Technology Assessment No. 112. AHRQ Publication No. 05-E009-2. Rockville MD. AHRQ. 2005.
- Vogel A, Hutchinson BL, Mitchell EA. Mastitis in the first year postpartum. *Birth*. 1999;26:218-225.
- Volante E, Moretti S, Pisani F, Bevilacqua G. Early diagnosis of bacterial infection in the neonate. *J Mat Fetal Neonat Med*. 2004;16 Suppl 2:13-6.
- Volmanen P, Valanne J, Alahuhta S. Breast-feeding problems after epidural analgesia for labour: a retrospective cohort study of pain, obstetrical procedures and breast-feeding practices. *Int J Obstet Anesth*. 2004;13(1):25-9.
- Watts DH, Lambert J, Stiehm ER, Harris DR, Bethel J, Mofenson L, Meyer WA 3rd, Mathieson B, Fowler MG, Nemo G, PACTG 185 Study Team. Progression of HIV disease among women following delivery. *J Acquir Immune Defic Syndr*. 2003;33(5):585-93.
- Weijers RN, Bekedam DJ, Smulders YM. Determinants of mild gestational hyperglycemia and gestational diabetes mellitus in a large dutch multiethnic cohort. *Diabetes Care*. 2002;25(1):72-7.
- Williams KP, Galerneau F. Intrapartum fetal heart rate patterns in the prediction of neonatal acedemia. *Am J Obstet Gynecol*. 2003;188: 820-3.
- Williams KP, Farquharson DF, Bebbington M, Dansereau J, Galerneau F, Wilson RD, Shaw D, Kent N. Screening for fetal well-being in a high-risk pregnant population comparing the nonstress test with umbilical artery Doppler velocimetry: a randomized controlled clinical trial. *Am J Obstet Gynecol*. 2003;188(5):1366-71.
- Wisborg K, Kesmodel U, Henriksen TB, Olsen SF, Secher NJ. Exposure to tobacco smoke in utero and the risk of stillbirth and death in the first year of life. *Am J Epidemiol*. 2001;154(4):322-7 .
- Wolfe EL, Davis T, Guldish J, Delucchi KL. Mortality risk associated with perinatal drug and alcohol use in California. *J Perinatol*. 2005;25(2):93-100.
- Wooley RJ. Benefits and risks of episiotomy: a review of English language literature since 1980. *Obstet Gynecol Surv*. 1995;50(11):806-820.
- Wong SF, Chow KM, Ho LC. The relative risk of 'fetal distress' in pregnancy associated meconium-stained liquor at different gestation. *J Obstet Gynaecol*. 2002; 22:594-9.

Wong C, Scavone B, Dugan S, Smith J, Prather H, Ganchiff J, McCarthy R. Incidence of postpartum lumbosacral spine and lower extremity nerve injuries. *Obstet Gynecol.* 2003;101:279-288.

World Health Organization Expert Consultation. Appropriate body mass index for Asian populations and its implication for policy and intervention strategies. *The Lancet.* 2004;363:157-163.

World Health Organization. Make every mother and child count. WHO Geneva. 2005

World Health Organization. Making a difference in countries. Strategic approach to improving maternal and newborn survival and health. WHO Geneva. 2006

Wu S, Kocherginsky M, Hibbard JU. Abnormal placentation: twenty year analysis. *Am J Obstet Gynecol.* 2005;192(5):1458-61.

Xiao R, Sorensen TK, Williams MA, Luthy DA. Influence of pre-eclampsia on fetal growth. *J Matern Fetal Neonatal Med.* 2003;13(3):157-62.

Xiong X, Buekens P, Alexander S, Demianczuk N, Wollast E. Anemia during pregnancy and birth outcome: a meta-analysis. *Am J Perinatol.* 2000;17(3):137-46.

Yang X, Hsu-Hage B, Zhang H, Zhang C, Zhang Y, Zhang C. Women with impaired glucose tolerance during pregnancy have significantly poorer pregnancy outcomes. *Diabetes Care.* 2002;25(9):1619-24.

Yogev Y, Langer O, Xenakis EM, Rosenn B. The association between glucose challenge test, obesity and pregnancy outcome in 6390 non-diabetic women. *J Matern Fetal Neonatal Med.* 2005;17(1):29-34.

Yokoe DS, Christiansen CL, Johnson R, Sands KE, Livingston J, Shtatland ES, Platt R. Epidemiology of and surveillance for postpartum infections. *Emerg Infect Dis.* 2001;7:837-841.

Yokoe DS, Noskin GA, Cunningham SM, Zuccotti G, Plaskett T, Fraser VJ, Olsen MA, Tokars JI, Solomon S, Perl TM, Cosgrove SE, Tilson RS, Greenbaum M, Hooper DC, Sands KE, Tully J, Herwaldt L, Diekema DJ, Wong ES, Climo M, Platt R. Enhanced identification of postoperative infections among inpatients. *Emerg Infect Dis.* 2004;10(11):1924-30.

Youssef R, Ramalingam U, Macleod M, Murphy DJ. Cohort study of maternal and neonatal morbidity in relation to use of episiotomy at instrumental vaginal delivery. *Br J Obstet Gynaecol.* 2005;112(7):941-5.

Zeitlin JA, Saurel-Cubizolles MJ, Ancel PY. *Paediatr Perinat Epidemiol.* 2002;6(2):124-30.

Zetterstrom K, Lindeberg SN, Haglund B, Hanson U. Maternal complications in women with chronic hypertension: a population-based cohort study. *Acta Obstet Gynecol Scand.* 2005;84(5):419-24.

Ziadeh SM, Sunna E. Obstetric and perinatal outcome of pregnancies with term labour and meconium-stained amniotic fluid. *Arch Gynecol Obstet.* 2000;264:84-7.

Zhang J, Troendle JF, Levine RJ. Risks of hypertensive disorders in the second pregnancy. *Paediatr Perinat Epidemiol.* 2001 Jul;15(3):226-31

Zhu BP, Haines KM, Le T, McGrath-Miller K, Boulton ML. Effect of the interval between pregnancies on perinatal outcomes among white and black women. *Am J Obstet Gynecol.* 2001;185(6):1403-10.

Zhu BP, Le T. Effect of interpregnancy interval on infant low birth weight: a retrospective cohort study using the Michigan Maternally Linked Birth Database. *Matern Child Health J.* 2003;7(3):169-78.

Zhu BP, Rolfs RT, Nangle BE, Horan JM. Effect of the interval between pregnancies on perinatal outcomes. *N Engl J Med.* 1999;340(8):589-94.